

Projekt na zachování
populací vzácných rostlin
bezlesého prostředí
v Čechách a v Sasku

UNIVERZITA J. E. PURKYNĚ V ÚSTÍ NAD LABEM



<http://biologie.ujep.cz/>



Ziel 3 | Cíl 3

Ahoj sousede. Hallo Nachbar.

2007-2013. www.ziel3-cil3.eu



Europäische Union. Europäischer Fonds für
regionale Entwicklung: Investition in Ihre
Zukunft / Evropská unie. Evropský fond pro
regionální rozvoj: Investice do vaší budoucnosti

Projekt- Nr./č. 100044665

Erhaltungsprojekt für
seltene Pflanzen
des Offenlandes
in Böhmen und Sachsen



**TECHNISCHE
UNIVERSITÄT
DRESDEN**

www.tu-dresden.de/bio/epobs

BIBLIOGRAPHIE

AUSZUG AUS DER
PROJEKTBEZOGENEN
LITERATURDATENBANK

INHALTSVERZEICHNIS

Antennaria dioica.....	3
Calamagrostis phragmitoides.....	10
Carex pulicaris.....	12
Carlina acaulis subsp. caulescens (subsp. simplex).....	16
Dactylorhiza sambucina.....	17
Dianthus seguieri (D. sylvaticus).....	19
Eriophorum latifolium.....	20
Gentianella amarella subsp. amarella.....	21
Gentianella germanica.....	21
Gentianella lutescens subsp. lutescens.....	21
Gentianella obtusifolia subsp. sturmiana.....	21
Hypochaeris maculata.....	25
Laserpitium prutenicum.....	26
Orchis mascula.....	27
Orchis morio.....	33
Orchis ustulata.....	41
Pedicularis palustris.....	42
Phyteuma orbiculare.....	43
Polygala serpyllifolia.....	44
Pseudorchis albida.....	45
Swertia perennis.....	47
Trifolium montanum.....	48
GesamtÜbersicht.....	51

ANTENNARIA DIOICA

- (1) Abadie, J. C.; Puttsepp, U.; Gebauer, G.; Faccio, A.; Bonfante, P. & Selosse, M. A. (2006), 'Cephalanthera longifolia (Neottieae, Orchidaceae) is mixotrophic: a comparative study between green and nonphotosynthetic individuals', *Canadian Journal of Botany-revue Canadienne De Botanique* **84**(9), 1462--1477.
- (2) Ahmed, A. A.; Hussein, T. A.; Mahmoud, A. A.; Farag, M. A.; Pare, P. W.; Wojcinska, M.; Karchesy, J. & Mabry, T. J. (2004), 'Nor-ent-kaurane diterpenes and hydroxylactones from *Antennaria geyeri* and *Anaphalis margaritacea*', *Phytochemistry* **65**(18), 2539--2543.
- (3) Aksenova, A. A.; Onipchenko, V. G. & Blinnikov, M. S. (1998), 'Plant interactions in alpine tundra: 13 years of experimental removal of dominant species', *Ecoscience* **5**(2), 258--270.
- (4) Andersson, U.-B. (2004), 'Botany day in Orebo county 2004', *Svensk Botanisk Tidskrift* **98**(5), 263--267.
- (5) Baltisberger, M. & Widmer, A. (2009), 'Karyological data of some angiosperms from Romania', *Willdenowia* **39**(2), 353--363.
- (6) Bayer, R. J. & Stebbins, G. L. (1987), 'Chromosome Numbers, Patterns of Distribution, and Apomixis in *Antennaria* (Asteraceae: Inuleae)', *Systematic Botany* **12**(2), 305-319.
- (7) Bayer, R. J. (1984), 'Chromosome Numbers and Taxonomic Notes for North American Species of *Antennaria* (Asteraceae: Inuleae)', *Systematic Botany* **9**(1), 74-83.
- (8) Bayer, R. J. (1989), 'A Taxonomic Revision of the *Antennaria rosea* (Asteraceae: Inuleae: Gnaphaliinae) Polyploid Complex', *Brittonia* **41**(1), 53-60.
- (9) Bayer, R. J. (1993), 'A Taxonomic Revision of the Genus *Antennaria* (Asteraceae: Inuleae: Gnaphaliinae) of Alaska and Yukon Territory, Northwestern North America', *Arctic and Alpine Research* **25**(2), 150-159.
- (10) Bayer, R. J.; Soltis, D. E. & Soltis, P. S. (1996), 'Phylogenetic Inferences in *Antennaria* (Asteraceae: Gnaphalieae: Cassiniinae) Based on Sequences from Nuclear Ribosomal DNA Internal Transcribed Spacers (ITS)', *American Journal of Botany* **83**(4), 516-527.
- (11) BICCHI, C.; NANO, G. M. & TIRA, S. (1975), 'Normal-paraffin Components of Some Gnaphalieae', *Planta Medica* **28**(4), 389--391.
- (12) Blachnik, T. (2009), 'Arnika und Katzenpfötchen - Ein Artenhilfsprojekt im Bayerischen Vogtland', Herausgeber:Regierung von Oberfranken, keine VA.
- (13) Blachnik, T. (2010), 'Artenhilfsprojekt Arnika und Katzenpfötchen im Bayerischen Vogtland, Landkreis Hof - Fachgrundlagen - Verbreitung, Bestandssituation, Erhaltungsmaßnahmen und Öffentlichkeitsarbeit für Leitarten bodensaurer

- Magerrasen', Technical report, Agentur und Naturschutzbüro Blachnik, keine VA, allgemeine Verbreitung Bayern/Vogtland.
- (14) Blinnikov, M. S. (1994), 'Experimental investigation of alpine plant communities in the Northwestern Caucasus: Phytolith analysis and holocene dynamics of alpine vegetation', Veroeffentlichungen des Geobotanischen Institutes der Eidgenoessische Technische Hochschule Stiftung Ruebel in Zuerich 0(115), 23--40.
- (15) Bobbink, R. & Roelofs, J. G. (1995), 'Ecological effects of atmospheric deposition on non-forest ecosystems in Western Europe', Studies in Environmental Science 64, 279 - 292.
- (16) Bohlin, K. (2006), 'Different colour forms in *Antennaria dioica*', Svensk Botanisk Tidskrift 100(1), 3--4.
- (17) BROWN, L. S. R. & GRAY, D. O. (1988), '1-amino-2-propanol, A Natural Product From the Compositae', Phytochemistry 27(4), 1195--1197.
- (18) BUHL, O.; FALCK, P.; KARSHOLT, O.; LARSEN, K. & SCHNACK, K. (1989), 'Records of Microlepidoptera From Denmark In 1987 Lepidoptera', Entomologiske Meddelelser 57(3), 123--135.
- (19) Chiryat'ev, E. A. & Rusakova, O. A. (1994), 'Species of Siberian flora as sources of direct action anti-coagulation', Rastitel'nye Resursy 30(4), 21--28.
- (20) Chmielewski, J. G. (1998), '*Antennaria dioica* (Asteraceae: Inuleae): Addition to the vascular flora of California', Madrono 45(3), 271--272.
- (21) CHMIELEWSKI, J. G.; CHINNAPPA, C. C. & SEMPLE, J. C. (1990), 'The Genus *Antennaria* (asteraceae, Inuleae) In Western North-america - Morphometric Analysis of *Antennaria-alborosea*, *Antennaria-corymbosa*, *Antennaria-marginata*, *Antennaria-microphylla*, *Antennaria-parvifolia*, *Antennaria-rosea*, and *Antennaria-umbrinella*', Plant Systematics and Evolution 169(3-4), 151--175.
- (22) Cranston, D. M. & Valentine, D. H. (1983), 'Transplant experiments on rare plant species from Upper Teesdale', Biological Conservation 26(2), 175 - 191.
- (23) CRAVEN, A. F. & CRAVEN, S. M. (1977), 'A 2nd Station For *Diapensia-lapponica* New-record', Watsonia 11(4), 376.
- (24) DAVYDOVA, M. V. (1982), 'Effect of Factory Smoke On Grassy Pine Forests In the Southern Urals Russian-sfsr Ussr', Biologicheskie Nauki (Moscow)(11), 71--74.
- (25) Dragana, S.; Comic, L.; Stefanovic, O. & Solujic-Sukdolak, S. (2008), 'In vitro antibacterial activity of *Antennaria dioica* L. and evaluation of synergism between plant extract and preservatives', Planta Medica 74(9), 1206--1206.
- (26) DUECK, T. A.; VANDEREERDEN, L. J. & BERDOWSKI, J. J. M. (1992), 'Estimation of So2 Effect Thresholds For Heathland Species', Functional Ecology 6(3), 291--296.

- (27) Eriksson, O. & Kiviniemi, K. (1999), 'Site occupancy, recruitment and extinction thresholds in grassland plants: an experimental study', *Biological Conservation* 87(3), 319 - 325.
- (28) Eriksson, O. (1996), 'Population ecology and conservation: Some theoretical considerations with examples from the Nordic flora', *Acta Universitatis Upsaliensis Symbolae Botanicae Upsalienses* 31(3), 159--167.
- (29) Eriksson, O. (1997), 'Colonization dynamics and relative abundance of three plant species (*Antennaria dioica*, *Hieracum pilosella* and *Hypochoeris maculata*) in dry semi-natural grasslands', *Ecography* 20(6), 559 - 568.
- (30) GABRIELIAN, E. T. & GAMBARYAN, P. P. (1973), 'New and Rare Floristic Finds In the Armenian-ssr', *Biologicheskii Zhurnal Armenii* 26(11), 56--60.
- (31) Heckford, R. J. (2001), 'Coleophora pappiferella Hofmann, 1869 (Lepidoptera: Coleophoridae) new to Britain and *Scrobipalpa murinella* (Duponchel, 1843) (Lepidoptera: Gelechiidae) at the same locality', *Entomologist's Gazette* 52(2), 101--104.
- (32) Heckford, R. J. (2001), 'Observations on the larval habits of *Kessleria saxifragae* (Stainton) (Lepidoptera: Yponomeutidae)', *Entomologist's Gazette* 52(1), 49--50.
- (33) Heckford, R. J. (2010), '*Scrobipalpa murinella* (Duponchel, 1843) (Lepidoptera: Gelechiidae) in Northern Ireland, and a note on the larva', *Entomologist's Gazette* 61(2), 132.
- (34) Heggland, A. (2005), 'New findings of lanceleaf grapefern *Botrychium lanceolatum* in Tinn, Telemark', *Blyttia* 63(4), 213--215.
- (35) HEIJNE, B.; HOFSTRA, J. J.; HEIL, G. W.; VANDAM, D. & BOBBINK, R. (1992), 'Effect of the Air-pollution Component Ammonium-sulfate On the Vam Infection-rate of 3 Heathland Species', *Plant and Soil* 144(1), 1--12.
- (36) Hogstrom, S. (1995), '*Prunella laciniata* in Gotland, Sweden', *Svensk Botanisk Tidskrift* 89(6), 333--334.
- (37) Hrivnak, R. & Ujhazy, K. (2005), 'Changes of the mountain grassland vegetation after abandonment and colonization by Norway spruce', *Ekologia-bratislava* 24(3), 231--253.
- (38) JONASSON, S. & CALLAGHAN, T. V. (1992), 'Root Mechanical-properties Related To Disturbed and Stressed Habitats In the Arctic', *New Phytologist* 122(1), 179--186.
- (39) KAMYSHEV, N. S. (1979), 'Comparative Characteristics of Dry Meadows and Caespitose Herb Steppes', *Biologicheskie Nauki (Moscow)*(3), 62--66.
- (40) Klanderud, K. (2010), 'Species recruitment in alpine plant communities: the role of species interactions and productivity', *Journal of Ecology* 98(5), 1128--1133.

- (41) Kleijn, D.; Bekker, R. M.; Bobbink, R.; De Graaf, M. C. C. & Roelofs, J. G. M. (2008), 'In search for key biogeochemical factors affecting plant species persistence in heathland and acidic grasslands: a comparison of common and rare species', *Journal of Applied Ecology* 45(2), 680--687.
- (42) Klett, J. E.; Staats, D. & Rogoyski, M. (2005), 'Preemergence weed control in container-grown herbaceous perennials', *Hortscience* 40(4), 1024.
- (43) KROEZE, C.; PEGTEL, D. M. & BLOM, C. J. C. (1989), 'An Experimental Comparison of Aluminum and Manganese Susceptibility In *Antennaria-dioica*, *Arnica-montana*, *Viola-canina*, *Filago-minima* and *Deschampsia-flexuosa*', *Acta Botanica Neerlandica* 38(2), 165--172.
- (44) Kytöviita, M.-M.; Vestberg, M. & Tuomi, J. (2003), 'A test of mutual aid in common mycorrhizal networks: established vegetation negates benefit in seedlings', *Ecology* 84(4), 898-906.
- (45) LYUBARSKII, E. L.; POLUYANOVA, V. I. & SILINA, L. G. (1975), 'Seasonal Variations of the Morphological Structure of Community Populations of Mouse-ear Hawkweed and Common Cats-ear', *Biologicheskie Nauki (Moscow)* 18(10), 75--79.
- (46) Mericli, A. H. (1983), 'Constituents of *Antennaria dioica*', *Journal of Natural Products* 46(6), 941-941.
- (47) MINARIK, J. (1950), '*Antennaria dioica* Gartn., *Gnaphalium dioicum* L.', *Cas Cesk Lek* 63(9-12), 254--60.
- (48) Mortensen, L. M. (1993), 'Effects of ozone on growth of several subalpine plant species', *Norwegian Journal of Agricultural Sciences* 7(2), 129--138.
- (49) Mückschel, C. & Otte, A. (2003), 'Morphometric parameters: an approach for the indication of environmental conditions on calcareous grassland', *Agriculture, Ecosystems & Environment* 98(1-3), 213 - 225.
- (50) NIKOLAEVA, V. G. (1977), 'Plants Used By Ethnic Groups of the Ussr For Treating Diseases of the Liver and Bile Secretory System', *Rastitel'nye Resursy* 13(2), 396--403.
- (51) Norderhaug, A.; Bakkevik, B. & Skogen, A. (1997), '*Dactylorhiza sambucina*, a threatened species in Norway?', *Blyttia* 55(2), 73--86.
- (52) Oester, M.; Ask, K.; Roemermann, C.; Tackenberg, O. & Eriksson, O. (2009), 'Plant colonization of ex-arable fields from adjacent species-rich grasslands: The importance of dispersal vs. recruitment ability', *Agriculture Ecosystems & Environment* 130(3-4), 93--99.
- (53) Oklejewicz, K. (1997), '*Veronica gentianoides* (Scrophulariaceae): An interesting synanthropic species in the Polish flora', *Fragmenta Floristica et Geobotanica Series Polonica* 4(0), 377--378.

- (54) Oldhammer, B. (1994), 'History of fires in central and northwestern Dalarna', *Svensk Botanisk Tidskrift* 88(5), 259--266.
- (55) Onipchenko, V. G.; Blinnikov, M. S. & Semenova, G. V. (1994), 'Experimental investigation of alpine plant communities in the Northwestern Caucasus: Shading experiments in alpine grasslands', *Veroeffentlichungen des Geobotanischen Institutes der Eidgenoessische Technische Hochschule Stiftung Ruebel in Zuerich* 0(115), 89--99.
- (56) Onipchenko, V. G.; Makarov, M. I. & van der Maarel, E. (2001), 'Influence of alpine plants on soil nutrient concentrations in a monoculture experiment', *Folia Geobotanica* 36(3), 225--241.
- (57) Öster, M. & Eriksson, O. (2007), 'Sex ratio mediated pollen limitation in the dioecious herb *Antennaria dioica*', *Ecoscience* 14(3), 387-398.
- (58) Pavlov, V. N.; Onipchenko, V. G.; Aksenova, A. A.; Volkova, E. V.; Zueva, O. I. & Makarov, M. I. (1998), 'The role of competition in Alpine plant communities (the Northwestern Causacus): An experimental approach', *Zhurnal Obshchei Biologii* 59(5), 453--476.
- (59) PEGTEL, D. M. (1988), 'Germination In Declining and Common Herbaceous Plant-populations Co-occurring In An Acid Peaty Heathland', *Acta Botanica Neerlandica* 37(2), 215--223.
- (60) PEGTEL, D. M.; BLOM, C. & KROEZE, C. (1987), 'Aluminum and Manganese Affected Growth of Threatened Plant Species Occurring On Poorly Buffered Sandy Soils of Moderate Acidity', *International Botanical Congress Abstracts* 17, 392.
- (61) Punt, W. & Hoen, P. (2009), 'The Northwest European Pollen Flora, 70: Asteraceae – Asteroideae', *Review of Palaeobotany and Palynology* 157(1-2), 22 - 183.
- (62) S. P. Carter, J. P. & Slingsby, D. R. (1987), 'Soil and Vegetation of the Keen of Hamar Serpentine, Shetland', *Journal of Ecology* 75, No. 1, 21-42.
- (63) Santos-Gonzalez, J. C.; Finlay, R. D. & Tehler, A. (2007), 'Seasonal dynamics of arbuscular mycorrhizal fungal communities in roots in a seminatural grassland', *Applied and Environmental Microbiology* 73(17), 5613--5623.
- (64) SCHWABE, A. (1990), 'Syndynamic Processes In Nardo-callunetea Communities Changes In Fallow Land After Renewed Cattle Grazing and Life History of *Arnica montana* L', *Carolinea* 48, 45--68.
- (65) Schwabe-Braun, A. (1990), 'Veränderungen in montanen Borstgrasrasen durch Düngung und Brachlegung: *Antennaria dioica* und *Vaccinium vitis-idaea* als Indikatoren', *Tuexenia* 10, 295-310.
- (66) SIMACHEV, V. I. (1980), 'Biological Foundations of Protection of Rare Relict Species of the Higher Plants of Leningrad Oblast Russian-sfsr Ussr With Special Reference To *Pulsatilla-vernalis* *Viscaria-alpina* and *Oxytropis-sordida*', *Botanicheskii Zhurnal* (St.

- Petersburg) 65(5), 725--737.
- (67) Skuhrava, M.; Skuhravy, V. & Hellrigl, K. (2002), 'Gall midge fauna (Cecidomyiidae, Diptera) of South Tyrol (2): Gall midges in the National Park Stilfser Joch and in Val Badia in the Dolomite Alps.', *Gredleriana* 2, 103--136.
- (68) Soudzilovskaia, N. A. & Onipchenko, V. G. (2005), 'Experimental investigation of fertilization and irrigation effects on an alpine heath, northwestern Caucasus, Russia', *Arctic Antarctic and Alpine Research* 37(4), 602--610.
- (69) Soudzilovskaia, N. A.; Onipchenko, V. G.; Cornelissen, J. H. C. & Aerts, R. (2005), 'Biomass production, N : P ratio and nutrient limitation in a Caucasian alpine tundra plant community', *Journal of Vegetation Science* 16(4), 399--406.
- (70) Stetsyuk, N. O. (1995), 'The floristic findings in the lower course of the Vorskla River', *Ukrayins'kyi Botanichnyi Zhurnal* 52(5), 639--646.
- (71) STRID, A. (1979), 'Contribution To the Flora of Mount Kajmakcalan Voras Oros Northern Greece', *Annales Musei Goulandris* 4(RECD. 1979).
- (72) TIRA, S.; GALEFFI, C. & DELLEMON.EM (1969), 'Flavonoids of *Antennaria Dioica* Gaertn - A Mnr Study', *Annali Di Chimica* 59(3), 284--.
- (73) TRINAJSTIC, I. (1972), 'A Contribution To the Phytosociological Typological Studies of the Field Vegetation of the Croatian Mountain Slopes', *Poljoprivredna Znanstvena Smotra*(28), 95--101.
- (74) Uotila, P. & Ahti, T. (2009), 'Additions to the vascular flora of the Berezovye Islands (Koiviston saaret), Karelian Isthmus, Russia', *Memoranda Societatis pro Fauna et Flora Fennica* 85(2), 33--44.
- (75) URBANSKA, K. (1965), 'Badania Cytologiczne Nad Doswiadczalnymi Mieszancami Pomiedzy *Antennaria Carpatica* (wahlb) Bl Et Fing la *Dioica* (l) Gaertn - Cytological Studies In Experimental Hybrids Between *Antennaria Carpatica* (wahlb) Bl Et Fing and A *Dioica* (l) Gaertn', *Acta Biologica Cracoviensia Series Botanica* 8(1), 51--.
- (76) URBANSKA, K. (1969), 'Embryological Investigations In *Antennaria* Gaertn .v. Micro- and Macrosporogenesis In Natural Hybrids Between A *Carpatica* (wahlenb) Bluff Etlfingerh and A *Dioica* (l) Gaertn', *Acta Biologica Cracoviensia Series Botanica* 12(1-2), 21--.
- (77) Van Den Berg, L. J. L.; Dorland, E.; Vergeer, P.; Hart, M. A. C.; Bobbink, R. & Roelofs, J. G. M. (2005), 'Decline of acid-sensitive plant species in heathland can be attributed to ammonium toxicity in combination with low pH', *New Phytologist* 166(2), 551--564.
- (78) van den Berg, L. J. L.; Peters, C. J. H.; Ashmore, M. R. & Roelofs, J. G. M. (2008), 'Reduced nitrogen has a greater effect than oxidised nitrogen on dry heathland vegetation', *Environmental Pollution* 154(3), 359--369.

- (79) Varga, S. & Kytöviita, M.-M. (2008), 'Sex-specific responses to mycorrhiza in a dioecious species', *Am. J. Bot.* 95(10), 1225-1232.
- (80) Varga, S. & Kytöviita, M.-M. (2010), 'Interrelationships between mycorrhizal symbiosis, soil pH and plant sex modify the performance of *Antennaria dioica*', *Acta Oecologica* 36(3), 291 - 298.
- (81) VERA DE LA PUENTE, M. L. (1985), 'Chorology of Acidophilous Taxa In the Cantabrian Mountains Spain', *Revista de Biología de la Universidad de Oviedo* 3, 71--84.
- (82) Viciani, D. & Moggi, G. (1996), 'Notes on the high altitude flora at Pratomagno (Eastern Tuscany) and considerations on the effects of the methane gas pipeline', *Webbia* 51(1), 59--81.
- (83) VOLKOVICH, V. B. (1972), 'Sex Ratios and Growth Characteristics In Diclinous Individuals of *Antennaria-dioica*', *Botanicheskii Zhurnal (St. Petersburg)* 57(10), 1278-1286.
- (84) von Ubisch, G. (1936), 'Genetic Studies on the Nature of Hermaphroditic Plants in *Antennaria Dioica* (L) Gaertn', *Genetics* 21(3), 282–294.
- (85) WALKER, R. B.; CHOWDAPPA, P. & GESSEL, S. P. (1993), 'Major-element Deficiencies In *Casuarina-equisetifolia*', *Fertilizer Research* 34(2), 127--133.
- (86) Westergaard, M. (1958), 'The Mechanism of Sex Determination in Dioecious Flowering Plants', *Advances in Genetics* 9, 217 - 281.
- (87) Wilhalm, T. (2007), 'Vascular plants', *Gredleriana* 7, 428--431.

CALAMAGROSTIS PHRAGMITOIDES

- (88) Brezina, S.; Koubek, T.; Münzbergová, Z. & Herben, T. (2006), 'Ecological benefits of integration of *Calamagrostis epigejos* ramets under field conditions', *Flora* 201(6), 461-467.
- (89) de Langhe, J.-E. & Delvosalle, L. (1973), 'Calamagrostis-phragmitoides An Indigenous Grass of Belgium', *Bulletin de la Societe Royale de Botanique de Belgique* 106(1), 61-64.
- (90) Dersch, G. & Mast, R. (2000), 'Distribution and community formation by *Calamagrostis phragmitoides* HARTMAN (Scandinavian Small-reed, Poaceae) in the German Harz Mountains', *Tuexenia* 20(20), 119-129.
- (91) Frey, L. & Paszko, B. (1999), 'Remarks on the distribution, taxonomy and karyology of *Calamagrostis* species (Poaceae) with special reference to their representatives in Poland', *Fragmenta floristica et geobotanica suppl.* 7, 33-45.
- (92) Gerstlauer, L. (1940), 'Eine umstrittene sächsische *Calamagrostis*.', *Hercynia* 3, 174-176.
- (93) Hackel, E. (1883), 'Über das Vorkommen von *Calamagrostis phragmitoides* (HARTM.) in Deutschland', *Bericht Deutsche Botanische Gesellschaft* 1, 161-162.
- (94) Hackel, E. (1883), 'Über das Vorkommen von *Calamagrostis phragmitioides* (HARTM.) in Deutschland', *Bericht Deutsche Botanische Gesellschaft* 1, 161-162.
- (95) Heine, R. (1970), 'Zur Taxonomie und Verbreitung von *Calamagrostis canescens* (WEB.) ROTH, *Calamagrostis villosa* (CHAIX) J. F.GMELIN, *Calamagrostis arundinacea* (L.) ROTH, *Calamagrostis neglecta* (EHRH.) PAL. BEAUV., *Calamagrostis pseudopurpurea* (GERSTL.) HEINE und einiger Bastarde in Sachsen (Bezirke Dresden, Karl-Marx-Stadt und Leipzig).', PhD thesis, TU Dresden, Fakultät für Bau-, Wasser- und Forstwesen.
- (96) Heine, R. (1971), '*Calamagrostis neglecta* (Ehrh.) Pal. Beauv. in der Oberlausitz', *Abhandlungen und Berichte des Naturkunde-Museum Görlitz* 46, 1-8.
- (97) Heine, R. (1972), 'Zur Klärung der sächsischen *Calamagrostis*-Art in den Muldentälern, dem Flöha- und dem Zschopautal.', *Feddes Repertorium* 83, 275-288.
- (98) Krause, A. & Wolf, G. (1991), 'Das Purpureitgras (*Calamagrostis purpurea* (Trin.) Trin. ssp. *phragmitoides* (Hartm.) Tzvelev) neu für Nordrhein-Westfalen', *Dechenia* 144, 115.
- (99) Ludwig, W. (1991), 'Purple Reed Grass *Calamagrostis-phragmitoides* In Hesse and Adjacent Regions', *Hessische Floristische Briefe* 40(2), 17-20.
- (100) Meinunger, L. (1977), 'Purpureitgras, *Calamagrostis phragmitoides* Hartman, in

- Thuringen und Sachsen gefunden.', *Hercynia* 14, 53-54.
- (101) Nieschalk, A. & Nieschalk, C. (1976), 'Nachtrag und Berichtigung zu einem Beitrag über *Calamagrostis phragmitoides* Hartman in Nordhessen', *Hessische Floristische Briefe* 25, 68-69.
- (102) Philippi, G. (1970), 'Calamagrostis phragmitoides (Hartm.), das Purpureitgras im Schwarzwald und in den Vogesen', *Beitraege zur Naturkundlichen Forschung in Suedwestdeutschland* 29(2), 107-110.
- (103) Rauschert, S. (1973), 'Der korrekte Name des Sachsichen Endemiten *Calamagrostis pseudopurpurea*', *Berichte der Arbeitsgemeinschaft Sächsischer Botaniker* 10, 25-29.
- (104) Rebele, F. & Lehmann, C. (2001), 'Biological flora of Central Europe: *Calamagrostis epigejos* (L.) Roth.', *Flora* 196, 325-344.
- (105) Schiebold, S. (2006), 'Zur Genetischen und Morphologischen Struktur, Reproduktionsökologie, Soziologischen Bindung und Verbreitung von *Calamagrostis Rivalis* H.Scholz im Einzugsgebiet der Mulde, Mitteldeutschland.', Master's thesis, Martin-Luther-Universität Halle-Wittenberg.
- (106) Schiebold, S.; Hensen, I.; Wesche, K. & Röser, M. (2009), 'Extensive clonality of the endemic *Calamagrostis pseudopurpurea* Gerstl. ex O. R. Heine in central Germany revealed by RAPD markers.', *Plant Biology* 11, 473-782.
- (107) Scholz, H. (1971), 'Gegenwärtige Kenntnisse über die Verbreitung der *Calamagrostis phragmitoides* Hartm. (Purpureitgras) in Mitteleuropa.', *Philippia* 1, 85-90.
- (108) Smejkal, M. (1976), 'Calamagrostis-phragmitoides New-record For the Czechoslovakian Flora', *Preslia (Prague)* 48(2), 124-130.

CAREX PULICARIS

- (109) Austrheim, G.; Gunilla, E.; Olsson, A. & Grøntvedt, E. (1999), 'Land-use impact on plant communities in semi-natural sub-alpine grasslands of Budalen, central Norway', *Biological Conservation* 87(3), 369 - 379.
- (110) Bartoszek, W. (1999), 'A new station for *Carex pulicaris* (Cyperaceae) in the Polish Carpathians', *Fragmenta Floristica et Geobotanica Series Polonica* 6(0), 274--277.
- (111) Bergamini, A.; Peintinger, M.; Fakheran, S.; Moradi, H.; Schmid, B. & Joshi, J. (2009), 'Loss of habitat specialists despite conservation management in fen remnants 1995-2006', *Perspectives in Plant Ecology, Evolution and Systematics* 11(1), 65 - 79.
- (112) Coroi, M.; Skeffington, M. S.; Giller, P.; Smith, C.; Gormally, M. & O'Donovan, G. (2004), 'Vegetation diversity and stand structure in streamside forests in the south of Ireland', *Forest Ecology and Management* 202(1-3), 39 - 57.
- (113) Dunnett, N.; Nagase, A.; Booth, R. & Grime, P. (2008), 'Influence of vegetation composition on runoff in two simulated green roof experiments', *Urban Ecosystems* 11(4), 385--398.
- (114) Escudero, M.; Valcárcel, V.; Vargas, P. & Luceño, M. (2010), 'Bipolar disjunctions in *Carex*: Long-distance dispersal, vicariance, or parallel evolution?', *Flora - Morphology, Distribution, Functional Ecology of Plants* 205(2), 118 - 127.
- (115) Eysink, F. & De Bruijn, O. (1997), 'Restoration of fen meadows in the Lemselermaten', *Levende Natuur* 98(7), 258--265.
- (116) Fyfe, R. M.; Brown, A. G. & Rippon, S. J. (2004), 'Characterising the late prehistoric, []Romano-British' and medieval landscape, and dating the emergence of a regionally distinct agricultural system in South West Britain', *Journal of Archaeological Science* 31(12), 1699 - 1714.
- (117) Garcia-Baquero Moneo, G. & Valle Gutierrez, C. J. (1999), 'New plant species in the Sierra de la Demanda (La Rioja, Spain)', *Studia Botanica* 18, 145--148.
- (118) Gorissen, I. (1992), 'Noteworthy occurrences of *Platanthera bifolia*, *Azolla filliculoides*, and other species at the southern edge of the Lower Rhine Basin', *Floristische Rundbriefe* 26(1), 21--22.
- (119) GROOTJANS, A. P. (1979), 'Some Remarks On the Relation Between Nitrogen Mineralization, Groundwater Table and Standing Crop In Wet Meadows', *Acta Botanica Neerlandica* 28(2-3), 234--235.
- (120) Hájek, M.; Horsák, M.; Hájková, P. & Díte, D. (2006), 'Habitat diversity of central European fens in relation to environmental gradients and an effort to standardise fen terminology in ecological studies', *Perspectives in Plant Ecology, Evolution and Systematics* 8(2), 97 - 114.

- (121) Jankovská, V. & Rybníček, K. (1988), 'The genus *Carex* in the late glacial and holocene of Czechoslovakia', *Aquatic Botany* 30(1-2), 23 - 37.
- (122) Jansen, A. J. M.; Grootjans, A. P. & Jalink, M. H. (2000), 'Hydrology of Dutch *Cirsio-Molinietum* meadows: Prospects for restoration', *Applied Vegetation Science* 3(1), 51--64.
- (123) Jurzyk, S. & Wrobel, M. (2003), 'Co-occurrence of two species *Molinia caerulea* L. and "red-list" species *Carex pulicaris* L. in Western Pomerania (Poland)', *Polish Journal of Ecology* 51(3), 363--367.
- (124) Leck, M. A. & Schütz, W. (2005), 'Regeneration of Cyperaceae, with particular reference to seed ecology and seed banks', *Perspectives in Plant Ecology, Evolution and Systematics* 7(2), 95 - 133.
- (125) Leps, J. (2004), 'What do the biodiversity experiments tell us about consequences of plant species loss in the real world?', *Basic and Applied Ecology* 5(6), 529 - 534.
- (126) Llamas, F.; Acedo, C.; Andres, J.; Carbo, R.; Perez, M. A. & Penas, A. (1993), 'Anatomy of the European species of *Carex* L. Sect. *Unciniiformes* Kuk. (Cyperaceae)', *Anales del Jardin Botanico de Madrid* 51(1), 97--109.
- (127) Mueller, J. & Kuhbier, H. (2006), 'History, development and vegetation dynamics of a small calcareous fen in the conservation area Reithbruch near Bremen (NW-Germany).', *Abhandlungen Naturwissenschaftlichen Verein zu Bremen* 46(1), 101--124.
- (128) MUELLER, W. & SCHUMACHER, W. (1986), 'The Distribution of Rare Species of the Calcareous Flat Moor *Caricion-davallianae* In Eifel West Germany', *Decheniana* 139, 200--201.
- (129) Müller, F. (2001), 'Verbreitung, Soziologie und Gefährdung von *Carex pulicaris* in Sachsen', *Ber. AGsB NF* 18, 35-48.
- (130) Netien, G. (1994), 'Addendum to the flora of Lyonnais: *Carex buxbaumii* Wahl. and *Carex hartmanii* Cajander', *Bulletin Mensuel de la Societe Linneenne de Lyon* 63(7), 267--268.
- (131) NIEMI, A. (1970), 'Occurrences of *Carex-pulicaris*-m On the Hango Peninsula South Finland', *Memoranda Societatis pro Fauna et Flora Fennica* 46, 39--42.
- (132) Olsson, K.-A. (2006), 'New endangered vascular plants in Skane', *Botaniska Notiser* 139(3), 22--33.
- (133) Pedersen, H. A. (1997), 'Two unnoticed occurrences of *Selaginella selaginoides*', *Flora og Fauna* 103(1), 17--18.
- (134) Peintinger, M. & Bergamini, A. (2006), 'Community structure and diversity of bryophytes and vascular plants in abandoned fen meadows', *Plant Ecology* 185(1), 1--

- 17.
- (135) Phoenix, G. K.; Johnson, D.; Grime, J. P. & Booth, R. E. (2008), 'Sustaining ecosystem services in ancient limestone grassland: importance of major component plants and community composition', *Journal of Ecology* 96(5), 894--902.
- (136) PLACKOWSKI, R. (1986), 'New Localities of Rare Species of Plants Near Radomsko In the Province of Piotrkow Trybunalski Central Poland', *Fragmenta Floristica et Geobotanica* 31-32(1-2), 45--48.
- (137) Plackowski, R. (2009), 'PHYTOSOCIOLOGICAL COMPARATIVE ANALYSIS OF TWO LOCALITIES OF CAREX PULICARIS L. IN THE ŁÓDŹ PROVINCE (CENTRAL POLAND)', *TEKA Komisji Ochrony i Kształtowania Środowiska Przyrodniczego* 6, 223--227.
- (138) S. P. Carter, J. P. & Slingsby, D. R. (1987), 'Soil and Vegetation of the Keen of Hamar Serpentine, Shetland', *Journal of Ecology* 75, No. 1, 21-42.
- (139) SCHUMACKER, R.; DIRKSE, G. & DE ZUTTERE, P. (1974), 'Carex-dioica New-record and Carex-pulicaris In the Nature Reserve Lands of the Hohe Venn Mountains Province of Liege Belgium', *Lejeunia*(74), 1-21.
- (140) STANOVA, V. & GRULICH, V. (1993), 'Floristical and Phytogeographical Characteristic of Rudava River Alluvium', *Biologia* 48(4), 407--410.
- (141) Topic, J. & Ilijanic, L. (2001), 'Carex pulicaris L. (Cyperaceae), a new species in Croatian flora', *Natura Croatica* 10(4), 423--426.
- (142) Trivedi, M. R.; Morecroft, M. D.; Berry, P. M. & Dawson, T. P. (2008), 'Potential effects of climate change on plant communities in three montane nature reserves in Scotland, UK', *Biological Conservation* 141(6), 1665 - 1675.
- (143) van der Linden, M.; Vickery, E.; Charman, D. J.; Borekens, P. & van Geel, B. (2008), 'Vegetation history and human impact during the last 300 years recorded in a German peat deposit', *Review of Palaeobotany and Palynology* 152(3-4), 158--175.
- (144) van der Linden, M.; Vickery, E.; Charman, D. J.; Broekens, P. & van Geel, B. (2008), 'Vegetation history and human impact during the last 300 years recorded in a German peat deposit', *Review of Palaeobotany and Palynology* 152(3-4), 158 - 175.
- (145) Wallnofer, B. (2008), 'On the alleged presence of Carex pulicaris (Cyperaceae) in Croatia', *Acta Botanica Croatica* 67(2), 245--246.
- (146) Weeda, E. J. & Keulen, S. M. A. (2007), 'Changes in the vegetation of the kathager beemden reserve, west of heerlen', *Natuurhistorisch Maandblad* 96(2), 21--29.
- (147) WITTIG, B. (1987), 'A Fountain Peat-plant Community With Carex-pulicaris Near Verden West Germany', *Abhandlungen Naturwissenschaftlichen Verein zu Bremen* 40(4), 395--402.
- (148) Wittig, B. (1996), 'Preserving and revitalization of vegetation from a headwater slope

- by grazing. First results', *Abhandlungen Naturwissenschaftlichen Verein zu Bremen* 43(2), 309--316.
- (149) Witting, B. (1998), 'Comparison of recent and historical records on *Caricion fuscae* (acid small-sedge fen) in the Elbe-Weser region', *Drosera* 98(2), 85--102.
- (150) ZALUSKI, T. (1978), 'Rare Plants of the Pilica Tributary Area Poland', *Acta Universitatis Nicolai Copernici Biologia* 22, 133--136.

CARLINA ACAULIS SUBSP. CAULESCENS (SUBSP. SIMPLEX)

- (151) Becker, U.; Colling, G.; Dostal, P.; Jakobsson, A. & Matthies, D. (2006), 'Local adaptation in the monocarpic perennial *Carlina vulgaris* at different spatial scales across Europe', *Oecologia* 150(3), 506-518.
- (152) Hempel, W. (1982), 'Über das Indigenat von *Carlina acaulis* am Pöhlberg bei Annaberg', *Gled* 9, 201-203.
- (153) Klinkhamer, P. G. L.; De Jong, T. J. & Meelis, E. (1991), 'The control of flowering in the monocarpic perennial *Carlina vulgaris*', *Oikos* 61(1), 88-95.
- (154) Kruse, J. (1987), 'Morphological Investigations of Involucral and Receptacular Bracts of the Genus *Carlina* L', *Flora* 179(1), 17-33.
- (155) Neumann, S. M. (1999), 'Populationsbiologische und autökologische Untersuchungen an *Carlina acaulis* ssp. *caulescens* (LAM.) Schlüßer et Martens 1834 im geographischen Vergleich', Master's thesis, Universität Göttingen.
- (156) Nickstadt, A. (2000), 'Populationsbiologie und ökologische Morphologie der Halbtrockenrasenart *Carlina acaulis* L.', Master's thesis, Universität Halle-Wittenberg.
- (157) Nickstadt, A. & Jäger, E. J. (2000), 'Beiträge zur Populationsbiologie der Silberdistel (*Carlina acaulis* L.)', *Hercynia* 33, 245-256.
- (158) Tauscher, L. (1978), 'Zum Vorkommen von *Carlina acaulis* L. subsp. *simplex* (W. et. K.) Arcang. auf dem Pöhlberg bei Annaberg (Westerzgebirge).', *Gleditschia* 6, 141-143.
- (159) Trejgell, A.; Bednarek, M. & Tretyn, A. (2009), 'Micropropagation of *Carlina acaulis*', *Acta Biologica Cracoviensia* 51, 97-103.
- (160) Trejgell, A.; Dabrowska, G. & Tretyn, A. (2009), 'In vitro regeneration of *Carlina acaulis* subsp. *simplex* from seedling explants', *Acta Physiologiae Plantarum* 31(3), 445-453.
- (161) Trejgell, A. & Tretyn, A. (2007), 'Analysis of flowering ability of regenerated *Carlina acaulis* subsp. *simplex* plants', *ACTA AGROBOTANICA* 60, 39-44.

DACTYLORHIZA SAMBUCINA

- (162) Gigord, L. D. B.; Macnair, M. R. & Smithson, A. (2001), 'Negative frequency-dependent selection maintains a dramatic flower color polymorphism in the rewardless orchid *Dactylorhiza sambucina* (L.) Soo', PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA 98, 6253-6255.
- (163) Inghe, O. & Tamm, C. O. (1988), 'Survival and flowering of perennial herbs. V. Patterns of flowering', Oikos 51(2), 203-219.
- (164) Jersakova, J.; Johnson, S. D.; Kindlmann, P. & Pupin, A. C. (2008), 'Effect of nectar supplementation on male and female components of pollination success in the deceptive orchid *Dactylorhiza sambucina*', Acta Oecologica 33(3), 300-306.
- (165) Jersakova, J.; Kindlmann, P. & Renner, S. S. (2006), 'Is the colour dimorphism in *Dactylorhiza sambucina* maintained by differential seed viability instead of frequency-dependent selection?', Folia Geobotanica 41(1), 61-76.
- (166) Juillet, N.; Dunand-Martin, S. & Gigord, L. D. B. (2007), 'Evidence for inbreeding depression in the food-deceptive colour-dimorphic orchid *Dactylorhiza sambucina* (L.) Soo', Plant Biology 9(1), 147-151.
- (167) Kropf, M. & Renner, S. S. (2005), 'Pollination success in monochromic yellow populations of the rewardless orchid *Dactylorhiza sambucina*', Plant Systematics and Evolution 254(3-4), 185-197.
- (168) Kropf, M. (1995), 'Vegetationskundlicher Vergleich von Standorten des Holunder-Knabenkrautes (*Dactylorhiza sambucina*) im Nahegebiet und der Rheinhessischen Schweiz unter Berücksichtigung von Pflege sowie Verbreitung und Gefährdung dieser Orchideenart.', Flora Fauna Rheinland-Pfalz 8, 133-145.
- (169) Mattiason, G. (1986), 'The Decline of *Dactylorhiza-sambucina* In the Nature Reserve Kungsmarken Southernmost Sweden', Svensk Botanisk Tidskrift 80(1), 17-25.
- (170) Nilsson, L. A. (1980), 'The Pollination Ecology of *Dactylorhiza-sambucina* (orchidaceae)', Botaniska Notiser 133(3), 367-385.
- (171) Nordal, I. & Wischmann, F. (1987), '*Dactylorhiza sambucina* in Norway [norwegisch]', Blyttia 45(1), 30-38.
- (172) Norderhaug, A.; Bakkevik, B. & Skogen, A. (1997), '*Dactylorhiza sambucina*, a threatened species in Norway? [norwegisch]', Blyttia 55(2), 73-86.
- (173) Pedersen, H. A. E. (2006), 'Systematics and evolution of the *Dactylorhiza romana/sambucina* polyploid complex (Orchidaceae)', Botanical Journal of the Linnean Society 152(4), 405-434.
- (174) Pellegrino, G. & Bellusi, F. (2009), 'Molecular identification of mycorrhizal fungi in

- Dactylorhiza sambucina* (Orchidaceae)', *Biologia* 64, 893-897.
- (175) Pellegrino, G. & Musacchio, A. (2006), 'Effects of defoliation on reproductive success in two orchids, *Serapias vomeracea* and *Dactylorhiza sambucina*', *Annales Botanici Fennici* 43(2), 123-128.
- (176) Pellegrino, G.; Bellusci, F. & Musacchio, A. (2005), 'Evidence of post-pollination barriers among three colour morphs of the deceptive orchid *Dactylorhiza sambucina* (L.) Soo', *Sexual Plant Reproduction* 18(4), 179-185.
- (177) Pellegrino, G.; Bellusci, F. & Musacchio, A. (2008), 'Double floral mimicry and the magnet species effect in dimorphic co-flowering species, the deceptive orchid *Dactylorhiza sambucina* and rewarding *Viola aethnensis*', *Preslia* 80(4), 411-422.
- (178) Pellegrino, G.; Caimi, D.; Noce, M. E. & Musacchio, A. (2005), 'Effects of local density and flower colour polymorphism on pollination and reproduction in the rewardless orchid *Dactylorhiza sambucina* (L.) Soo', *Plant Systematics and Evolution* 251(2-4), 119-129.
- (179) Pettersson, G. & Nilsson, L. A. (1983), 'The Pollination Ecology of *Dactylorhiza-sambucina* On the Island of Stora Karlsö Sweden In the Baltic', *Svensk Botanisk Tidskrift* 77(2), 123-132.
- (180) Walentoswki, H. & Obermeier, E. (1992), 'Mountain meadows with *Dactylorhiza sambucina* (L.) Soo at the Brotjacklriegel in the Vorderer Bayerischer Wald (Bavaria)', *Tuexenia* 0(12), 193-208.

DIANTHUS SEGUIERI (D. SYLVATICUS)

- (181) Beinhauer, H. (2001), 'Populationsgenetische Untersuchungen an Vorkommen der seltenen Pflanzenart *Dianthus seguieri* VILL. in Mittel-und Ostdeutschland unter Verwendung eines molekulargenetischen Markersystems (RAPD)', PhD thesis, Friedrich-Schiller-Universität.
- (182) Zibareva, L.; Volodin, V.; Saatov, Z.; Savchenko, T.; Whiting, P.; Lafont, R. & Dinan, L. (2003), 'Distribution of phytoecdysteroids in the Caryophyllaceae', *Phytochemistry* 64(2), 499 - 517.

ERIOPHORUM LATIFOLIUM

- (183) Gruber, M. (1984), 'The Marshes With Eriophorum latifolium and Sedges In the Valley of Louron Central Pyrenees France', Bulletin de la Societe d'Histoire Naturelle de Toulouse 120, 27-30.
- (184) Phillips, M. E. (1954), 'Eriophorum Angustifolium Roth (e Polystachion L)', Journal of Ecology 42(2), 612-622.
- (185) Turland, N. (1997), 'Proposal to reject the name Eriophorum polystachion (Cyperaceae)', Taxon 46(1), 115-116.
- (186) Vitko, K. R. (1981), 'Ecology of Cotton-grass Eriophorum-latifolium (Cyperaceae) In the Moldavian-SSR', Izvestiya Akademii Nauk Moldavskoi SSR Seriya Biologicheskikh i Khimicheskikh Nauk 2(2), 30-32.

GENTIANELLA AMARELLA SUBSP. AMARELLA

GENTIANELLA GERMANICA

GENTIANELLA LUSTESCENS SUBSP. LUSTESCENS

GENTIANELLA OBTUSIFOLIA SUBSP. STURMIANA

- (187) Akhalkatsi, M. & Wagner, J. (1996), 'Reproductive phenology and seed development of *Gentianella caucasea* in different habitats in the central caucasus', *Flora* 191(2), 161-168.
- (188) Brunzel, S. (2010), 'Ex-Situ Cultivation and In-Situ Management as Contribution to the Protection of Species - The examples of *Cypripedium calceolus*, *Gladiolus palustris*, *Gentianella bohemica* and *Gentianella lutescens*', *Naturschutz und Landschaftsplanung* 42(5), 148-156.
- (189) Dierschke, H. (1986), 'Untersuchungen zur Populationsdynamik der *Gentianella*-Arten in einem Enzian.Zwenken-Kalkmagerrasen', *Natur und Heimat* 46, 73-81.
- (190) Eriksen, M.; Bjureke, K. E. & Dhillion, S. S. (2002), 'Mycorrhizal plants of traditionally managed boreal grasslands in Norway', *Mycorrhiza* 12(3), 117-123.
- (191) Fischer, M. & Matthies, D. (1997), 'Mating structure and inbreeding and outbreeding depression in the rare plant *Gentianella germanica* (Gentianaceae)', *American Journal of Botany* 84(12), 1685-1692.
- (192) Fischer, M. & Matthies, D. (1997), 'The effect of population size on reproduction, population growth rate and offspring fitness in the rare plant *Gentianella germanica*', *Bulletin of the Ecological Society of America* 78(4 SUPPL.), 247.
- (193) Fischer, M. & Matthies, D. (1998), 'Effects of population size on performance in the rare plant *Gentianella germanica*', *Journal of Ecology* 86(2), 195-204.
- (194) Fischer, M. & Matthies, D. (1998), 'Experimental demography of the rare *Gentianella germanica*: seed bank formation and microsite effects on seedling establishment', *Ecography* 21(3), 269-278.
- (195) Fischer, M. & Matthies, D. (1998), 'RAPD variation in relation to population size and plant fitness in the rare *Gentianella germanica* (Gentianaceae)', *American Journal of Botany* 85(6), 811-819.
- (196) Fischer, M.; Matthies, D. & Schmid, B. (1997), 'Responses of rare calcareous grassland plants to elevated CO₂: a field experiment with *Gentianella germanica* and *Gentiana cruciata*', *Journal of Ecology* 85(5), 681-691.

- (197) Gargano, D.; Bellusci, F.; Pellegrino, G.; Palermo, A. M.; Bernardo, L. & Musacchio, A. (2009), 'The conservation perspectives and value of small and isolated plant populations: preliminary clues for *Gentianella crispata* (Gentianaceae) at the western boundary of its range', *Annales Botanici Fennici* 46(2), 115-124.
- (198) Greimler, J. & Dobes, C. (2000), 'High genetic diversity and differentiation in relict lowland populations of *Gentianella austriaca* (A. and J. Kern.) Holub (Gentianaceae)', *Plant Biology* 2(6), 628-637.
- (199) Greimler, J.; Hermanowski, B. & Jang, C. G. (2004), 'A re-evaluation of morphological characters in European *Gentianella* section *Gentianella* (Gentianaceae)', *Plant Systematics and Evolution* 248(1-4), 143-169.
- (200) Grzes, I. M. (2007), 'Does rare *Gentianella germanica* (Wild.) Borner originating from calamine spoils differ in selected morphological traits from reference populations?', *Plant Species Biology* 22(1), 49-52.
- (201) Hempel, W. (1980), 'Die Verbreitung der Kleinenziane in Sachsen', *Naturschutzarbeit naturkundliche Heimatforschung Sachsen* 22, 36-44.
- (202) Hempel, W. (1981), 'Die sächsischen Sippen des Formenkreises von *Gentianella germanica* (Willd.) Borner', *Gleditschia* 8, 31-41.
- (203) Holub, J. (1967), 'Neue Namen innerhalb der Gattungen *Gentianella* Moench, *Gentianopsis* Ma und *Comastoma* (Wettst.) Tokoyuni', *Folia Geobotanica & Phytotaxonomica* 2, 115-120.
- (204) Huhta, A. P.; Hellstrom, K.; Rautio, P. & Tuomi, J. (2003), 'Grazing tolerance of *Gentianella amarella* and other monocarpic herbs: why is tolerance highest at low damage levels?', *Plant Ecology* 166(1), 49-61.
- (205) Huhta, A. P.; Lennartsson, T.; Tuomi, J.; Rautio, P. & Laine, K. (2000), 'Tolerance of *Gentianella campestris* in relation to damage intensity: an interplay between apical dominance and herbivory', *Evolutionary Ecology* 14(4-6), 373-392.
- (206) Jang, C. G.; Muellner, A. N. & Greimler, J. (2005), 'Conflicting patterns of genetic and morphological variation in European *Gentianella* section *Gentianella*', *Botanical Journal of the Linnean Society* 148(2), 175-187.
- (207) Juenger, T.; Lennartsson, T. & Tuomi, J. (2000), 'The evolution of tolerance to damage in *Gentianella campestris*: natural selection and the quantitative genetics of tolerance', *Evolutionary Ecology* 14(4-6), 393-419.
- (208) Kelly, D. (1989), 'Demography of Short-lived Plants In Chalk Grassland: 3. Population Stability', *Journal of Ecology* 77(3), 785-798.
- (209) Kelly, D. (1989), 'Demography of Short-lived Plants In Chalk Grassland: 2. Control of Mortality and Fecundity', *Journal of Ecology* 77(3), 770-784.

- (210) Kelly, D. (1989), 'Demography of Short-lived Plants In Chalk Grassland: 1. Life-cycle Variation In Annuals and Strict Biennials', *Journal of Ecology* 77(3), 747-769.
- (211) Kery, M. & Matthies, D. (2004), 'Reduced fecundity in small populations of the rare plant *Gentianopsis ciliate* (Gentianaceae)', *Plant Biology* 6(6), 683-688.
- (212) Kesting, S. (2001), 'Systematik, Beiträge zur Biologie und Naturschutzmanagement des *Gentianella campestris* agg. (l.) Börner', Master's thesis, Universität Halle-Wittenberg.
- (213) Kirschnerova, L. & Kirschner, J. (2003), 'Zmeny v rozsireni *Gentianella lutescens* subsp. *lutescens* a subsp. *carpatica* Ceske republice. [Changes in the distribution of *Gentianella lutescens* subsp. *lutescens* and subsp. *carpatica* in the Czech Republic.]', *Zprávy České Botanické Společnosti* 38, 205-216.
- (214) Krumbiegel, A. (2008), 'Socialization of *Gentianella campestris* ssp. *baltica* at the east bank of the Murtitz Lake (Mecklenburg-Western Pomerania) and near Ballenstedt at the north rim of the Harz Mountains (Saxony-Anhalt).', *Hercynia* 41(2), 219-238.
- (215) Lennartsson, T. & Oostermeijer, J. G. B. (2001), 'Demographic variation and population viability in *Gentianella campestris*: Effects of grassland management and environmental stochasticity', *Journal of Ecology* 89(3), 451-463.
- (216) Lennartsson, T. & Svensson, R. (1996), 'Patterns in the decline of three species of *Gentianella* (Gentianaceae) in Sweden, illustrating the deterioration of semi-natural grasslands', *Acta Universitatis Upsaliensis Symbolae Botanicae Upsalienses* 31(3), 169-184.
- (217) Lennartsson, T. (1996), 'Om gentianornas gronings- och blomningsfenologi [Phenology of germination and flowering in *Gentianella*]', *Svensk Botanisk Tidskrift* 90(5-6), 263-265.
- (218) Lennartsson, T. (1997), 'Seasonal differentiation - a conservative reproductive barrier in two grassland *Gentianella* (Gentianaceae) species', *Plant Systematics and Evolution* 208(1-2), 45-69.
- (219) Lennartsson, T.; Oostermeijer, J. G. B.; van Dijk, J. & den Nijs, H. C. M. (2000), 'Ecological significance and heritability of floral reproductive traits in *Gentianella campestris* (Gentianaceae).', *Basic and Applied Ecology* 1(1), 69-81.
- (220) Luijten, S. H.; Oostermeijer, J. G. B.; Ellis-Adam, A. C. & Den Nijs, H. C. M. (1998), 'Reproductive biology of the rare biennial *Gentianella germanica* compared with other gentians of different life history', *Acta Botanica Neerlandica* 47(3), 325-336.
- (221) McVeigh, A.; Carey, J. E. & Rich, T. C. G. (2005), 'Chiltern Gentian, *Gentianella germanica* (Willd.) Börner (Gentianaceae) in Britain: distribution and current status', *Watsonia* 25(Part 4), 339-367.
- (222) Milberg, P. (1994), 'Germination Ecology of the Endangered Grassland Biennial

- Gentianella campestris*', *Biological Conservation* 70(3), 287-290.
- (223) Oostermeijer, J. G. B.; Luijten, S. H.; Ellis-Adam, A. C. & den Nijs, J. C. M. (2002), 'Future prospects for the rare, late-flowering *Gentianella germanica* and *Gentianopsis ciliata* in Dutch nutrient-poor calcareous grasslands', *Biological Conservation* 104(3), 339-350.
- (224) Paland, S. & Schmid, B. (2003), 'Population size and the nature of genetic load in *Gentianella germanica*', *Evolution* 57(10), 2242-2251.
- (225) Petanidou, T.; Dennijs, J. C. M. & Oostermeijer, J. G. B. (1995), 'Pollination Ecology and Constraints On Seed Set of the Rare Perennial *Gentiana-cruciata* L. In the Netherlands', *Acta Botanica Neerlandica* 44(1), 55-74.
- (226) Petanidou, T.; Ellis-Adam, A. C.; Den Nijs, J. C. M. & Oostermeijer, J. G. B. (1998), 'Pollination ecology of *Gentianella uliginosa*, a rare annual of the Dutch coastal dunes', *Nordic Journal of Botany* 18(5), 537-548.
- (227) Prekorsek, B. (1972), 'Contribution To the Problems of Pseudoseasonal Polymorphism of *Gentianella-praecox*', *Bioloski Vestnik*(20), 17-29.
- (228) Rich, T.; Holyoak, D.; Margetts, L. & Murphy, R. (1997), 'Hybridisation between *Gentianella amarella* and *G. anglica*', *Watsonia* 21, 313-325.
- (229) Skalicky, V. (1969), 'The Collective Species *Gentianella germanica* in the Bohemian Massiv', *Preslia* 41(2), 140-147.
- (230) Stadler, K.; Koch, M.; Bernhardt, K. G. & Greimler, J. (2010), 'Spatial arrangement and genetic structure in *Gentianella aspera* in a regional, local, and temporal context', *Plant Systematics and Evolution* 286(1-2), 7-19.
- (231) Wagner, J. & Mitterhofer, E. (1998), 'Phenology, seed development, and reproductive success of an alpine population of *Gentianella germanica* in climatically varying years', *Botanica Acta* 111(2), 159-166.
- (232) Winfield, M. O.; Wilson, P. J.; Labra, M. & Parker, J. S. (2003), 'A brief evolutionary excursion comes to an end: the genetic relationship of British species of *Gentianella* sect. *Gentianella* (*Gentianaceae*)', *Plant Systematics and Evolution* 237(3-4), 137-151.

HYPOCHAERIS MACULATA

- (233) Eriksson, O. (1997), 'Colonization dynamics and relative abundance of three plant species (*Antennaria dioica*, *Hieracium pilosella* and *Hypochaeris maculata*) in dry semi-natural grasslands', *Ecography* 20(6), 559-568.
- (234) Öster, M.; Persson, K. & Eriksson, O. (2008), 'Validation of plant diversity indicators in semi-natural grasslands', *Agriculture, Ecosystems & Environment* 125(1-4), 65 - 72.
- (235) Öster, M.; Ask, K.; Cousins, S. A. O. & Eriksson, O. (2009), 'Dispersal and establishment limitation reduces the potential for successful restoration of semi-natural grassland communities on former arable fields', *Journal of Applied Ecology* 46(6), 1266-1274.
- (236) Wells, T. C. E. (1976), 'Biological Flora of the British Isles: *Hypochaeris maculata*', *Journal of Ecology* 64(2), 757-774.

LASERPITIUM PRUTENICUM

- (237) Bohlmann, F. & Zdero, C. (1971), 'Terpene derivatives from higher plants .11. prutenin, 4-angeloyloxypruteninone, and 4-acetoxypruteninone and 4-acetoxyisopruteninone - 4 new sesquiterpene lactons from Laserpitium-prutenicum l', chemische Berichte-recueil 104(5), 1611-.
- (238) RYCHLEWSKA, U.; HODGSON, D.; HOLUB, M.; BUDESINSKY, M. & SMITALOVA, Z. (1985), 'ON TERPENES .288. THE STRUCTURE OF 2-OXO-8-ALPHA-ANGELOYLOXY-11-ALPHA-ACETOXY-5-BETA-H,6-ALPHA-H,7-ALPHA-H- GUAI-1(10),3-DIEN-6,12-OLIDE, A SESQUITERPENE LACTONE FROM LASERPITIUM-PRUTENICUM L REVISION OF THE STEREOSTRUCTURES OF NATIVE 2-OXOGUAI-1(10),3-DIEN-6,12-OLIDES FROM THE SPECIES OF THE UMBELLIFERAE FAMILYS', COLLECTION OF CZECHOSLOVAK CHEMICAL COMMUNICATIONS 50(11), 2607-2624.

ORCHIS MASCULA

- (239) Aceto, S.; Montieri, S.; Sica, M. & Gaudio, L. (2007), 'Molecular evolution of the OrcPI locus in natural populations of Mediterranean orchids', *Gene* 392(1-2), 299 - 305.
- (240) Adriaens, D.; Honnay, O. & Hermy, M. (2006), 'No evidence of a plant extinction debt in highly fragmented calcareous grasslands in Belgium', *Biological Conservation* 133(2), 212 - 224.
- (241) Alpaslan, M. & Hayta, M. (2007), 'Effect of soymilk substitution on the rheological and sensory properties of salep (Traditional turkish milk beverage)', *INTERNATIONAL JOURNAL OF FOOD PROPERTIES* 10(3), 413-420.
- (242) Atalla, R.; Hackney, J.; Uhlin, I. & Thompson, N. (1993), 'Hemicelluloses as structure regulators in the aggregation of native cellulose', *International Journal of Biological Macromolecules* 15(2), 109 - 112.
- (243) Aybeke, M.; Sezik, E. & Olgun, G. (2010), 'Vegetative anatomy of some Ophrys, Orchis and Dactylorhiza (Orchidaceae) taxa in Trakya region of Turkey', *Flora - Morphology, Distribution, Functional Ecology of Plants* 205(2), 73 - 89.
- (244) Aziz, N.; Mehmood, M. H.; Siddiqi, H. S.; Mandukhail, S.-u.-R.; Sadiq, F.; Maan, W. & Gilani, A. H. (2009), 'Antihypertensive, antidyslipidemic and endothelial modulating activities of *Orchis mascula*', *HYPERTENSION RESEARCH* 32(11), 997-1003.
- (245) Barkham, J. (1992), 'The effects of management on the ground flora of ancient woodland, Brigsteer Park Wood, Cumbria, England', *Biological Conservation* 60(3), 167 - 187.
- (246) Baskin, C. C. & Baskin, J. M. (1998), *Germination Ecology of Plants with Specialized Life Cycles and/or Habitats*'Seeds', Academic Press, San Diego, pp. 459 - 557.
- (247) Breadmore, K. N. & Kirk, W. D. J. (1998), 'Factors affecting floral herbivory in a limestone grassland', *Acta Oecologica* 19(6), 501 - 506.
- (248) Cafasso, D.; Pellegrino, G.; Caputo, P.; Scrugli, A. & Cozzolino, S. (2001), 'Genetic relatedness of insular segregates of mediterranean orchid species as inferred from ITS sequences analysis', *BOTANICA HELVETICA* 111(2), 181-190.
- (249) Calster, H. V.; Endels, P.; Antonio, K.; Verheyen, K. & Hermy, M. (2008), 'Coppice management effects on experimentally established populations of three herbaceous layer woodland species', *Biological Conservation* 141(10), 2641 - 2652.
- (250) Cantone, C.; Sica, M.; Gaudio, L. & Aceto, S. (2009), 'The OrcPI locus: Genomic organization, expression pattern, and noncoding regions variability in *Orchis italica* (Orchidaceae) and related species', *Gene* 434(1-2), 9 - 15.
- (251) Carruthers, W.; Duncan, F. & Shrubsole, W. (1875), 'Excursion to the isle of Sheppey:

- June 7th, 1875', *Proceedings of the Geologists' Association* 4(5), 320 - 323.
- (252) Cescutti, P.; Campa, C.; Delben, F. & Rizzo, R. (2002), 'Structure of the oligomers obtained by enzymatic hydrolysis of the glucomannan produced by the plant *Amorphophallus konjac*', *Carbohydrate Research* 337(24), 2505 - 2511.
- (253) Cescutti, P.; Pupulin, R.; Delben, F.; Abbate, M.; Dentini, M.; Sparapano, L.; Rizzo, R. & Crescenzi, V. (2002), 'New exopolysaccharides produced by *Aureobasidium pullulans* grown on glucosamine', *Carbohydrate Research* 337(13), 1203 - 1209.
- (254) Cozzolino, S. & Widmer, A. (2005), 'Orchid diversity: an evolutionary consequence of deception?', *Trends in Ecology & Evolution* 20(9), 487 - 494.
- (255) Cozzolino, S.; Aceto, S.; Caputo, P.; Gaudio, L. & Nazzaro, R. (1998), 'Phylogenetic relationships in *Orchis* and some related genera: an approach using chloroplast DNA', *NORDIC JOURNAL OF BOTANY* 18(1), 79-87.
- (256) Cozzolino, S.; Nardella, A.; Impagliazzo, S.; Widmer, A. & Lexer, C. (2006), 'Hybridization and conservation of Mediterranean orchids: Should we protect the orchid hybrids or the orchid hybrid zones?', *BIOLOGICAL CONSERVATION* 129(1), 14-23.
- (257) Cozzolino, S.; Schiestl, F.; Muller, A.; De Castro, O.; Nardella, A. & Widmer, A. (2005), 'Evidence for pollinator sharing in Mediterranean nectar-mimic orchids: absence of pre-mating barriers?', *PROCEEDINGS OF THE ROYAL SOCIETY B-BIOLOGICAL SCIENCES* 272(1569), 1271-1278.
- (258) D'Emerico, S.; Cozzolino, S.; Pellegrino, G.; Pignone, D. & Scrugli, A. (2002), 'Heterochromatin distribution in selected taxa of the 42-chromosomes *Orchis* s. l. (*Orchidaceae*)', *CARYOLOGIA* 55(1), 55-62.
- (259) Dormont, L.; Delle-Vedove, R.; Bessiere, J. M.; Hossaert-Mc Key, M. & Schatz, B. (2010), 'Rare white-flowered morphs increase the reproductive success of common purple morphs in a food-deceptive orchid', *NEW PHYTOLOGIST* 185(1), 300-310.
- (260) Honnay, O.; Hermy, M. & Coppin, P. (1999), 'Effects of area, age and diversity of forest patches in Belgium on plant species richness, and implications for conservation and reforestation', *Biological Conservation* 87(1), 73 - 84.
- (261) Honnay, O.; Hermy, M. & Coppin, P. (1999), 'Impact of habitat quality on forest plant species colonization', *Forest Ecology and Management* 115(2-3), 157 - 170.
- (262) Hopper, S. D. & Lambers, H. (2009), 'Darwin as a plant scientist: a Southern Hemisphere perspective', *Trends in Plant Science* 14(8), 421 - 435.
- (263) Hossain, M. M. (2010), 'Therapeutic Orchids: Traditional Uses and Recent Advances - An Overview', *Fitoterapia In Press, Accepted Manuscript*, - .
- (264) Inghe, O. & Tamm, C. O. (1988), 'Survival and Flowering of Perennial Herbs. V

- Patterns of Flowering', *Oikos* 51, 203-219.
- (265) Jacquemyn, H.; Brys, R.; Adriaens, D.; Honnay, O. & Roldan-Ruiz, I. (2009), 'Effects of population size and forest management on genetic diversity and structure of the tuberous orchid *Orchis mascula*', *CONSERVATION GENETICS* 10(1), 161-168.
- (266) Jacquemyn, H.; Brys, R.; Hermy, M. & Willems, J. H. (2005), 'Does nectar reward affect rarity and extinction probabilities of orchid species? An assessment using historical records from Belgium and the Netherlands', *Biological Conservation* 121(2), 257 - 263.
- (267) Jacquemyn, H.; Brys, R.; Honnay, O. & Hermy, M. (2008), 'Effects of coppicing on demographic structure, fruit and seed set in *Orchis mascula*', *BASIC AND APPLIED ECOLOGY* 9(4), 392-400.
- (268) Jacquemyn, H.; Brys, R.; Honnay, O. & Hutchings, M. J. (2009), 'Biological Flora of the British Isles: *Orchis mascula* (L.) L.', *Journal of Ecology* 97(2), 360--377.
- (269) Jacquemyn, H.; Honnay, O.; Cammue, B. P. A.; Brys, R. & Lievens, B. (2010), 'Low specificity and nested subset structure characterize mycorrhizal associations in five closely related species of the genus *Orchis*', *MOLECULAR ECOLOGY* 19(18), 4086-4095.
- (270) Jacquemyn, H.; Wiegand, T.; Vandepitte, K.; Brys, R.; Roldan-Ruiz, I. & Honnay, O. (2009), 'Multigenerational analysis of spatial structure in the terrestrial, food-deceptive orchid *Orchis mascula*', *Journal of Ecology* 97(2), 206--216.
- (271) Jersáková, J.; Johnson, S. D.; Kindlmann, P. & Pupin, A.-C. (2008), 'Effect of nectar supplementation on male and female components of pollination success in the deceptive orchid *Dactylorhiza sambucina*', *Acta Oecologica* 33(3), 300 - 306.
- (272) Johnson, M.; Putwain, P. & Holliday, R. (1978), 'Wildlife conservation value of derelict metalliferous mine workings in Wales', *Biological Conservation* 14(2), 131 - 148.
- (273) Johnson, S. & Nilsson, L. (1999), 'Pollen carryover, geitonogamy, and the evolution of deceptive pollination systems in orchids', *ECOLOGY* 80(8), 2607-2619.
- (274) Jones, E. & Hughes, R. E. (1983), 'Foliar ascorbic acid in some angiosperms', *Phytochemistry* 22(11), 2493 - 2499.
- (275) Knudsen, J. T.; Tollsten, L. & Bergström, L. (1993), 'Floral scents—a checklist of volatile compounds isolated by head-space techniques', *Phytochemistry* 33(2), 253 - 280.
- (276) Koch, M. A.; Scheriau, C.; Schupfner, M. & Bernhardt, K.-G. (2010), 'Long-term monitoring of the restoration and development of limestone grasslands in north western Germany: Vegetation screening and soil seed bank analysis', *Flora - Morphology, Distribution, Functional Ecology of Plants* In Press, Corrected Proof, - .
- (277) Kohn, D. (2005), 'The miraculous season', *NATURAL HISTORY* 114(9), 38-40.

- (278) Kull, T.; Kindlmann, P.; Hutchings, M. J. & Primack, R. B. (2006), 'Conservation biology of orchids: Introduction to the special issue', *Biological Conservation* 129(1), 1 - 3.
- (279) Larsson, E. & Örstadius, L. (2008), 'Fourteen coprophilous species of *Psathyrella* identified in the Nordic countries using morphology and nuclear rDNA sequence data', *Mycological Research* 112(10), 1165 - 1185.
- (280) Lievens, B.; van Kerckhove, S.; Justé, A.; Cammue, B. P.; Honnay, O. & Jacquemyn, H. (2010), 'From extensive clone libraries to comprehensive DNA arrays for the efficient and simultaneous detection and identification of orchid mycorrhizal fungi', *Journal of Microbiological Methods* 80(1), 76 - 85.
- (281) Majetic, C.; Rausher, M. & Raguso, R. (2010), 'The pigment-scent connection: Do mutations in regulatory vs. structural anthocyanin genes differentially alter floral scent production in *Ipomoea purpurea*?', *South African Journal of Botany* 76(4), 632 - 642.
- (282) Montieri, S.; Gaudio, L. & Aceto, S. (2004), 'Isolation of the LFY/FLO homologue in *Orchis italica* and evolutionary analysis in some European orchids', *Gene* 333, 101 - 109.
- (283) Nilsson, L. A. (1983), 'Anthecology of *Orchis mascula* (Orchidaceae)', *Nordic Journal of Botany* 3(2), 157--179.
- (284) Oklejewicz, K. (1997), '*Veronica gentianoides* (Scrophulariaceae): An interesting synanthropic species in the Polish flora', *Fragmenta Floristica et Geobotanica Series Polonica* 4(0), 377--378.
- (285) Pegg, G. & Vessey, J. (1973), 'Chitinase activity in *Lycopersicon esculentum* and its relationship to the in vivo lysis of *Verticillium albo-atrum* mycelium', *Physiological Plant Pathology* 3(2), 207 - 222.
- (286) Pellegrino, G.; Caputo, P.; Cozzolino, S.; Menale, B. & Musacchio, A. (2000), 'Molecular characterization of a hybrid zone between *Orchis mascula* and *O. pauciflora* in Southern Italy', *BIOLOGIA PLANTARUM* 43(1), 13-18.
- (287) Pellegrino, G.; Cozzolino, S.; D'Emérico, S. & Grunanger, P. (2000), 'The taxonomic position of the controversial taxon *Orchis clandestina* (Orchidaceae): Karyomorphological and molecular analyses', *BOTANICA HELVETICA* 110(2), 101-107.
- (288) Pellegrino, G.; D'Emérico, S.; Musacchio, A.; Scrugli, A. & Cozzolino, S. (2005), 'Confirmation of hybridization among sympatric insular populations of *Orchis mascula* and *O. provincialis*', *PLANT SYSTEMATICS AND EVOLUTION* 251(2-4), 131-142.
- (289) Peterken, G. (1974), 'A method for assessing woodland flora for conservation using indicator species', *Biological Conservation* 6(4), 239 - 245.
- (290) Peterken, G. (1977), 'Habitat conservation priorities in British and European

- woodlands', *Biological Conservation* 11(3), 223 - 236.
- (291) Peterken, G. F. & Francis, J. L. (1999), 'Open spaces as habitats for vascular ground flora species in the woods of central Lincolnshire, UK', *Biological Conservation* 91(1), 55 - 72.
- (292) Plowright, C. B. (1896), 'On the recent additions to our knowledge of the uredine[modifier letter high tone bar] and ustilagine[modifier letter high tone bar] with special reference to the british species', *Transactions of the British Mycological Society* 1, 90 - 105.
- (293) Poschlod, P. & WallisDeVries, M. F. (2002), 'The historical and socioeconomic perspective of calcareous grasslands—lessons from the distant and recent past', *Biological Conservation* 104(3), 361 - 376.
- (294) Rich, T. C. G. & Woodruff, E. R. (1996), 'Changes in the vascular plant floras of England and Scotland between 1930-1960 and 1987-1988: The BSBI Monitoring Scheme', *Biological Conservation* 75(3), 217 - 229.
- (295) Rosén, E. & Bakker, J. P. (2005), 'Effects of agri-environment schemes on scrub clearance, livestock grazing and plant diversity in a low-intensity farming system on Öland, Sweden', *Basic and Applied Ecology* 6(2), 195 - 204.
- (296) Salzmann, C. C.; Cozzolin, S. & Schiestl, F. P. (2007), 'Floral scent in food-deceptive orchids: Species specificity and sources of variability', *PLANT BIOLOGY* 9(6), 720-729.
- (297) Singh, R.; Kachroo, M. & Agrawal, S. S. (2009), 'Standardization of *Centipeda minima*, *Mimusops elengi*, *Orchis mascula* and *Paeonia emodii*', *INDIAN JOURNAL OF TRADITIONAL KNOWLEDGE* 8(3, Sp. Iss. SI), 338-345.
- (298) STEINBRUCK, G.; SCHLEGEL, M.; DAHLSTROM, I. & ROTTGER, B. (1986), 'CHARACTERIZATION OF INTERSPECIFIC HYBRIDS BETWEEN *ORCHIS-MASCULA* AND *ORCHIS-PALLENS* (ORCHIDACEAE) BY ENZYME ELECTROPHORESIS', *PLANT SYSTEMATICS AND EVOLUTION* 153(3-4), 229-241.
- (299) Tekinsen, K. K. & Güner, A. (2010), 'Chemical composition and physicochemical properties of tubera salep produced from some Orchidaceae species', *Food Chemistry* 121(2), 468 - 471.
- (300) Trist, P. (1981), '*Fritillaria meleagris* L.: Its survival and habitats in suffolk, England', *Biological Conservation* 20(1), 5 - 14.
- (301) Ülo Mander; Mikk, M. & Külvik, M. (1999), 'Ecological and low intensity agriculture as contributors to landscape and biological diversity', *Landscape and Urban Planning* 46(1-3), 169 - 177.
- (302) Ur-Rahman, A. & Zaman, K. (1989), 'Medicinal plants with hypoglycemic activity', *Journal of Ethnopharmacology* 26(1), 1 - 55.

- (303) Valletta, A.; Attorre, F.; Bruno, F. & Pasqua, G. (2008), 'In vitro asymbiotic germination of *Orchis mascula* L', *PLANT BIOSYSTEMS* 142(3), 653-655.
- (304) Vanderpoorten, A.; Delescaille, L.-M. & Jacquemart, A.-L. (2004), 'The bryophyte layer in a calcareous grassland after a decade of contrasting mowing regimes', *Biological Conservation* 117(1), 11 - 18.
- (305) von Oheimb, G. & Brunet, J. (2007), 'Dalby Söderskog revisited: long-term vegetation changes in a south Swedish deciduous forest', *Acta Oecologica* 31(2), 229 - 242.
- (306) Warcup, J. & Talbot, P. (1966), 'Perfect states of some rhizoctonias', *Transactions of the British Mycological Society* 49(3), 427 - 435.

ORCHIS MORIO

- (307) (1983), 'Abstracts of papers presented at the 20th Annual Meeting of the Society of Cryobiology August 30-September 2, 1983, University of Cambridge, Cambridge, United Kingdom Session 1–Symposium: Cryopreservation of organs', *Cryobiology* 20(6), 698 - 748.
- (308) (2006), Arzneipflanzenporträts, in Siegfried Bäuml, ed., 'Heilpflanzenpraxis Heute (First Edition)', Urban & Fischer, Munich, pp. 49 - 460.
- (309) (2008), 'SEB Main Meeting, Marseille 2008', *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology* 150(3, Supplement 1), Abstracts of the Annual Main Meeting of the Society of Experimental Biology, 6th - 10th July 2008, Marseille, France, keine VA, S1 - S40.
- (310) Aceto, S.; Cozzolino, S.; Gaudio, L.; Nazzaro, R. & De Luca, P. (1999), 'Pollination flow in hybrid formation between *Orchis morio* and *Orchis papilionacea* (Orchidaceae) in two different habitats', *INTERNATIONAL JOURNAL OF PLANT SCIENCES* 160(6), 1153-1156.
- (311) Alonso-Sande, M.; Teijeiro-Osorio, D.; Remuñán-López, C. & Alonso, M. (2009), 'Glucomannan, a promising polysaccharide for biopharmaceutical purposes', *European Journal of Pharmaceutics and Biopharmaceutics* 72(2), 453 - 462.
- (312) Barreto, D. W. & Parente, J. P. (2006), 'Chemical properties and biological activity of a polysaccharide from *Cyrtopodium cardiochilum*', *Carbohydrate Polymers* 64(2), 287 - 291.
- (313) Benincasa, M.-A.; Cartoni, G. & Fratte, C. D. (2002), 'Flow field-flow fractionation and characterization of ionic and neutral polysaccharides of vegetable and microbial origin', *Journal of Chromatography A* 967(2), 219 - 234.
- (314) BEYRL, H.; SMITH, S.; PETERSON, R. & FRANCO, C. (1995), 'COLONIZATION OF ORCHIS-MORIO PROTOCORMS BY A MYCORRHIZAL FUNGUS - EFFECTS OF NITROGEN NUTRITION AND GLYPHOSATE IN MODIFYING THE RESPONSES', *CANADIAN JOURNAL OF BOTANY-REVUE CANADIENNE DE BOTANIQUE* 73(8), 1128-1140.
- (315) Buchala, A. J.; Franz, G. & Meier, H. (1974), 'A glucomannan from the tubers of *Orchis morio*', *Phytochemistry* 13(1), 163 - 166.
- (316) Bullock, J. M. (1998), 'Community translocation in Britain: Setting objectives and measuring consequences', *Biological Conservation* 84(3), 199 - 214.
- (317) Cafasso, D.; Pellegrino, G.; Caputo, P.; Scrugli, A. & Cozzolino, S. (2001), 'Genetic relatedness of insular segregates of mediterranean orchid species as inferred from ITS sequences analysis', *BOTANICA HELVETICA* 111(2), 181-190.
- (318) Caputo, P.; Aceto, S.; Cozzolino, S. & Nazzaro, R. (1997), 'Morphological and

- molecular characterization of a natural hybrid between *Orchis laxiflora* and *O-morio* (Orchidaceae)', *PLANT SYSTEMATICS AND EVOLUTION* 205(3-4), 147-155.
- (319) Challe, J. & Struik, P. (2008), 'The impact on orchid species abundance of gathering their edible tubers by HIV/AIDS orphans: a case of three villages in the Southern Highlands of Tanzania', *NJAS - Wageningen Journal of Life Sciences* 56(3), 261 - 279.
- (320) Chrungoo, N. (1992), 'Concepts of dormancy regulation in vegetative plant propagules: A review', *Environmental and Experimental Botany* 32(4), 309 - 318.
- (321) Coates, F.; Lunt, I. D. & Tremblay, R. L. (2006), 'Effects of disturbance on population dynamics of the threatened orchid *Prasophyllum correctum* D.L. Jones and implications for grassland management in south-eastern Australia', *Biological Conservation* 129(1), 59 - 69.
- (322) Cozzolino, S.; Aceto, S.; Caputo, P.; Gaudio, L. & Nazzaro, R. (1998), 'Phylogenetic relationships in *Orchis* and some related genera: an approach using chloroplast DNA', *NORDIC JOURNAL OF BOTANY* 18(1), 79-87.
- (323) Cozzolino, S.; Schiestl, F.; Muller, A.; De Castro, O.; Nardella, A. & Widmer, A. (2005), 'Evidence for pollinator sharing in Mediterranean nectar-mimic orchids: absence of pre-mating barriers?', *PROCEEDINGS OF THE ROYAL SOCIETY B-BIOLOGICAL SCIENCES* 272(1569), 1271-1278.
- (324) Darkwa, A. & Scott, P. (2008), 'Growth and influence of defoliation in terrestrial orchids, *Anacamptis morio* and *Serapia lingua*', *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology* 150(3, Supplement 1), Abstracts of the Annual Main Meeting of the Society of Experimental Biology, 6th - 10th July 2008, Marseille, France, S197 - S197.
- (325) Darkwa, A. A. & Scott, P. (2009), 'Seasonal growth of temperate terrestrial orchids as affected by the old tuber', *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology* 153(2, Supplement 1), Abstracts of the Annual Main Meeting of the Society of Experimental Biology, 28th June - 1st July, Glasgow, UK, S212 - S212.
- (326) de Faria, A. L. (1997), 'The atomic basis of biological symmetry and periodicity', *Biosystems* 43(2), 115 - 135.
- (327) DEMERICO, S.; BIANCO, P. & MEDAGLI, P. (1993), 'CYTOLOGICAL AND KARYOLOGICAL STUDIES ON ORCHIDACEAE', *CARYOLOGIA* 46(4), 309-319.
- (328) DEMERICO, S.; Bianco, P. & Pignone, D. (1996), 'Cytomorphological characterization of diploid and triploid individuals of *Orchis x gennarii* Reichenb fil (Orchidaceae)', *CARYOLOGIA* 49(2), 153-161.
- (329) D'Emerico, S.; Galasso, I.; Pignone, D. & Scrugli, A. (2001), 'Localization of rDNA loci by fluorescent in situ hybridization in some wild orchids from Italy (Orchidaceae)',

- CARYOLOGIA 54(1), 31-36.
- (330) DEmerico, S.; Pignone, D. & Bianco, P. (1996), 'Karyomorphological analyses and heterochromatin characteristics disclose phyletic relationships among $2n=32$ and $2n=36$ species of *Orchis* (Orchidaceae)', *PLANT SYSTEMATICS AND EVOLUTION* 200(1-2), 111-124.
- (331) DIJK, E. & ECK, N. (1995), 'EFFECTS OF MYCORRHIZAL FUNGI ON IN-VITRO NITROGEN RESPONSE OF SOME DUTCH INDIGENOUS ORCHID SPECIES', *CANADIAN JOURNAL OF BOTANY-REVUE CANADIENNE DE BOTANIQUE* 73(8), 1203-1211.
- (332) Dijk, E. & Grootjans, A. (1998), 'Performance of four *Dactylorhiza* species over a complex trophic gradient', *ACTA BOTANICA NEERLANDICA* 47(3), 351-368.
- (333) Dijk, E. (1990), 'Effects of mycorrhizal fungi on in-vitro nitrogen response of juvenile orchids', *Agriculture, Ecosystems & Environment* 29(1-4), 91 - 97.
- (334) Essl, F. (2003), 'Remarkable floristic records from Vienna, Lower Austria, Burgenland and Styria.', *Linzer Biologische Beitrage* 35(2), 935--956.
- (335) Essl, F. (2005), 'Remarkable floristic records from Vienna, Lower Austria, Burgenland and Syria, Part II', *Linzer Biologische Beitrage* 37(2), 1207--1230.
- (336) Franz, G. (1973), 'Biosynthesis of salep mannan', *Phytochemistry* 12(10), 2369 - 2373.
- (337) Franz, G. (1979), 'Metabolism of Reserve Polysaccharides in Tubers of *Orchis morio* L', *Planta Med* 36(5), 68-73.
- (338) Gillman, M. & Dodd, M. (1998), 'The variability of orchid population size', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* 126(1-2), 65-74.
- (339) Gillman, M. P. & Dodd, M. (2000), 'Detection of delayed density dependence in an orchid population', *Journal of Ecology* 88(2), 204--212.
- (340) Grabherr, G. (2009), 'Biodiversity in the high ranges of the Alps: Ethnobotanical and climate change perspectives', *Global Environmental Change* 19(2), 167 - 172.
- (341) Grayer, R. J. & Kokubun, T. (2001), 'Plant-fungal interactions: the search for phytoalexins and other antifungal compounds from higher plants', *Phytochemistry* 56(3), 253 - 263.
- (342) Grubb, P. J. (1998), 'A reassessment of the strategies of plants which cope with shortages of resources', *Perspectives in Plant Ecology, Evolution and Systematics* 1(1), 3 - 31.
- (343) Grünanger, P.; Caporali, E.; Marziani, G.; Menguzzato, E. & Servettaz, O. (1998), 'Molecular (RAPD) analysis on Italian taxa of the *Ophrys bertolonii* aggregate (Orchidaceae)', *PLANT SYSTEMATICS AND EVOLUTION* 212(3-4), 177-184.
- (344) H. F. Beyrle, S. E. Smith, C. M. M. F. R. L. P. (1995), 'Colonization of *Orchis morio*

- protocorms by a mycorrhizal fungus: effects of nitrogen nutrition and glyphosate in modifying the responses', *Canadian Journal of Botany* 73, 1128-1140.
- (345) Haavik, S.; Paulsen, B. S.; Wold, J. K. & Øivind Grimme (1982), 'Arabinogalactans in a purified allergen preparation from pollen of timothy (*Phleum pratense*)', *Phytochemistry* 21(8), 1913 - 1919.
- (346) Hejcman, M.; Schellberg, J. & Pavlu, V. (2010), 'Dactylorhiza maculata, *Platanthera bifolia* and *Listera ovata* survive N application under P limitation', *Acta Oecologica* In Press, Corrected Proof, - .
- (347) Hodgson, J.; Grime, J.; Wilson, P.; Thompson, K. & Band, S. (2005), 'The impacts of agricultural change (1963-2003) on the grassland flora of Central England: processes and prospects', *Basic and Applied Ecology* 6(2), 107 - 118.
- (348) Hrevusová, Z.; Hejcman, M.; Pavlu, V. V.; Hakl, J.; Klaudivsová, M. & Mrkvicka, J. (2009), 'Long-term dynamics of biomass production, soil chemical properties and plant species composition of alluvial grassland after the cessation of fertilizer application in the Czech Republic', *Agriculture, Ecosystems & Environment* 130(3-4), 123 - 130.
- (349) Irving, A. (1875), 'The Geology of the Nottingham District.–Part II', *Proceedings of the Geologists' Association* 4(1), 57 - 90.
- (350) Jacquemyn, H.; Vandepitte, K.; Brys, R.; Honnay, O. & Roldan-Ruiz, I. (2007), 'Fitness variation and genetic diversity in small, remnant populations of the food deceptive orchid *Orchis purpurea*', *BIOLOGICAL CONSERVATION* 139(1-2), 203-210.
- (351) Janecek, S. & Leps, J. (2005), 'Effect of litter, leaf cover and cover of basal internodes of the dominant species *Molinia caerulea* on seedling recruitment and established vegetation', *Acta Oecologica* 28(2), 141 - 147.
- (352) Janecková, P.; Wotavová, K.; Schödelbauerová, I.; Jersáková, J. & Kindlmann, P. (2006), 'Relative effects of management and environmental conditions on performance and survival of populations of a terrestrial orchid, *Dactylorhiza majalis*', *Biological Conservation* 129(1), 40 - 49.
- (353) Jersakova, J. & Johnson, S. (2006), 'Lack of floral nectar reduces self-pollination in a fly-pollinated orchid', *OECOLOGIA* 147(1), 60-68.
- (354) Jersakova, J. & Kindlmann, P. (1998), 'Patterns of pollinator-generated fruit set in *Orchis morio* (Orchidaceae)', *FOLIA GEOBOTANICA* 33(4), 377-390.
- (355) Johnson, S. & Nilsson, L. (1999), 'Pollen carryover, geitonogamy, and the evolution of deceptive pollination systems in orchids', *ECOLOGY* 80(8), 2607-2619.
- (356) Johnson, S.; Peter, C. & Agren, J. (2004), 'The effects of nectar addition on pollen removal and geitonogamy in the non-rewarding orchid *Anacamptis morio*', *PROCEEDINGS OF THE ROYAL SOCIETY OF LONDON SERIES B-BIOLOGICAL SCIENCES*

- 271(1541), 803-809.
- (357) Johnson, S.; Peter, C.; Nilsson, L. & Agren, J. (2003), 'Pollination success in a deceptive orchid is enhanced by co-occurring rewarding magnet plants', *ECOLOGY* 84(11), 2919-2927.
- (358) Juillet, N. & Scopece, G. (2010), 'Does floral trait variability enhance reproductive success in deceptive orchids?', *Perspectives in Plant Ecology, Evolution and Systematics* 12(4), 317 - 322.
- (359) Juillet, N.; Dunand-Martin, S. & Gigord, L. D. B. (2007), 'Evidence for inbreeding depression in the food-deceptive colour-dimorphic orchid *Dactylorhiza sambucina* (L.) Soo', *PLANT BIOLOGY* 9(1), 147-151.
- (360) Kristiansen, K. A.; Freudenstein, J. V.; Rasmussen, F. N. & Rasmussen, H. N. (2004), 'Molecular identification of mycorrhizal fungi in *Neuwiedia veratrifolia* (Orchidaceae)', *Molecular Phylogenetics and Evolution* 33(2), 251 - 258.
- (361) Kropf, M. & Renner, S. S. (2008), 'Pollinator-mediated selfing in two deceptive orchids and a review of pollinium tracking studies addressing geitonogamy', *OECOLOGIA* 155(3), 497-508.
- (362) Landi, M.; Frignani, F.; Lazzeri, C. & Angiolini, C. (2009), 'Abundance of orchids on calcareous grasslands in relation to community species, environmental, and vegetational conditions', *RUSSIAN JOURNAL OF ECOLOGY* 40(7), 486-494.
- (363) Masuhara, G.; Katsuya, K. & Yamaguchi, K. (1993), 'Potential for symbiosis of *Rhizoctonia solani* and binucleate *Rhizoctonia* with seeds of *Spiranthes sinensis* var. *amoena* in vitro', *Mycological Research* 97(6), 746 - 752.
- (364) McKendrick, S. (1996), 'The effects of shade on seedlings of *Orchis morio* and *Dactylorhiza fuchsii* in chalk and clay soil', *NEW PHYTOLOGIST* 134(2), 343-352.
- (365) McKendrick, S. L. (1996), 'The effects of fertilizer and root competition on seedlings of *Orchis morio* and *Dactylorhiza fuchsii* in chalk and clay soil', *New Phytologist* 134(2), 335-342.
- (366) Menkovic, N.; Savikin, K.; Tasic, S.; Zdunic, G.; Stesevic, D.; Milosavljevic, S. & Vincek, D. (2010), 'Ethnobotanical study on traditional uses of wild medicinal plants in Prokletije Mountains (Montenegro)', *Journal of Ethnopharmacology* In Press, Corrected Proof, - .
- (367) Moccia, M. D.; Widmer, A. & Cozzolino, S. (2007), 'The strength of reproductive isolation in two hybridizing food-deceptive orchid species', *MOLECULAR ECOLOGY* 16(14), 2855-2866.
- (368) Montieri, S.; Gaudio, L. & Aceto, S. (2004), 'Isolation of the LFY/FLO homologue in *Orchis italica* and evolutionary analysis in some European orchids', *Gene* 333, 101 - 109.

- (369) NEILAND, M. & WILCOCK, C. (1995), 'MAXIMIZATION OF REPRODUCTIVE SUCCESS BY EUROPEAN ORCHIDACEAE UNDER CONDITIONS OF INFREQUENT POLLINATION', *PROTOPLASMA* 187(1-4), 39-48.
- (370) NILSSON, L. (1992), 'ANIMAL POLLINATORS ADJUST PLANT GENDER IN RELATION TO FLORAL DISPLAY - EVIDENCE FROM ORCHIS-MORIO (ORCHIDACEAE)', *EVOLUTIONARY TRENDS IN PLANTS* 6(1), 33-40.
- (371) Oostermeijer, J. G. B.; Luijten, S. H. & den Nijs, J. C. M. (2003), 'Integrating demographic and genetic approaches in plant conservation', *Biological Conservation* 113(3), 389 - 398.
- (372) Öztürk, M.; Çelik, A.; Güvensen, A. & Hamzaoglu, E. (2008), 'Ecology of tertiary relict endemic *Liquidambar orientalis* Mill. forests', *Forest Ecology and Management* 256(4), 510 - 518.
- (373) Pfeifer, M.; Passalacqua, N. G.; Bartram, S.; Schatz, B.; Croce, A.; Carey, P. D.; Kraudelt, H. & Jeltsch, F. (2010), 'Conservation priorities differ at opposing species borders of a European orchid', *Biological Conservation* 143(9), 2207 - 2220.
- (374) Ramsbottom, J. (1922), 'Orchid Mycorrhiza: With plates II-VII', *Transactions of the British Mycological Society* 8(1-2), 28 - 61, IN2-IN7.
- (375) ROSSI, W.; CORRIAS, B.; ARDUINO, P.; CIANCHI, R. & BULLINI, L. (1992), 'GENE VARIATION AND GENE FLOW IN ORCHIS-MORIO (ORCHIDACEAE) FROM ITALY', *PLANT SYSTEMATICS AND EVOLUTION* 179(1-2), 43-58.
- (376) Ryan, M. J. & Smith, D. (2004), 'Fungal genetic resource centres and the genomic challenge', *Mycological Research* 108(12), 1351 - 1362.
- (377) Schatz, B. (2006), 'Fine scale distribution of pollinator explains the occurrence of the natural orchid hybrid *X Orchis bergonii*', *ECOSCIENCE* 13(1), 111-118.
- (378) Schiestl, F. (2005), 'On the success of a swindle: pollination by deception in orchids', *NATURWISSENSCHAFTEN* 92(6), 255-264.
- (379) Schiestl, F. P.; Johnson, S. D. & Raguso, R. A. (2010), 'Floral evolution as a figment of the imagination of pollinators', *Trends in Ecology & Evolution* 25(7), 382 - 383.
- (380) Schlüter, P. M. & Schiestl, F. P. (2008), 'Molecular mechanisms of floral mimicry in orchids', *Trends in Plant Science* 13(5), 228 - 235.
- (381) Scopece, G.; Cozzolino, S.; Johnson, S. D. & Schiestl, F. P. (2010), 'Pollination Efficiency and the Evolution of Specialized Deceptive Pollination Systems', *AMERICAN NATURALIST* 175(1), 98-105.
- (382) Scopece, G.; Juillet, N.; Mueller, A.; Schiestl, F. P. & Cozzolino, S. (2009), 'Pollinator attraction in *Anacamptis papilionacea* (Orchidaceae): a food or a sex promise?', *PLANT SPECIES BIOLOGY* 24(2), 109-114.

- (383) Sharma, I. K.; Jones, D. L. & French, C. J. (2003), 'Unusually high genetic variability revealed through allozymic polymorphism of an endemic and endangered Australian orchid, *Pterostylis* aff. *picta* (Orchidaceae)', *Biochemical Systematics and Ecology* 31(5), 513 - 526.
- (384) Silvertown, J.; Wells, D. A.; Gillman, M.; Dodd, M. E.; Robertson, H. & Lakhani, K. H. (1994), 'Short-term effects and long-term after-effects of fertilizer application on the flowering population of green-winged orchid *Orchis morio*', *Biological Conservation* 69(2), 191 - 197.
- (385) Smith, S. E. & Read, D. J. (1997), *Orchid mycorrhizas* (Mycorrhizal Symbiosis (Second Edition)), Academic Press, London, pp. 349 - 375.
- (386) Smithson, A. (2002), 'The consequences of rewardlessness in orchids: Reward-supplementation experiments with *Anacamptis morio* (Orchidaceae)', *AMERICAN JOURNAL OF BOTANY* 89(10), 1579-1587.
- (387) Smithson, A. (2006), 'Pollinator limitation and inbreeding depression in orchid species with and without nectar rewards', *NEW PHYTOLOGIST* 169(2), 419-430.
- (388) Stambouli-Meziane, H.; Bouazza, M. & Thinon, M. (2009), 'La diversité floristique de la végétation psammophile de la région de Tlemcen (nord-ouest Algérie)', *Comptes Rendus Biologies* 332(8), 711 - 719.
- (389) Tekinsen, K. K. & Güner, A. (2010), 'Chemical composition and physicochemical properties of tubera salep produced from some Orchidaceae species', *Food Chemistry* 121(2), 468 - 471.
- (390) Tsavkelova, E. A.; Bömke, C.; Netrusov, A. I.; Weiner, J. & Tudzynski, B. (2008), 'Production of gibberellic acids by an orchid-associated *Fusarium proliferatum* strain', *Fungal Genetics and Biology* 45(10), 1393 - 1403.
- (391) Tsiftsis, S.; Tsiripidis, I.; Karagiannakidou, V. & Alifragis, D. (2008), 'Niche analysis and conservation of the orchids of east Macedonia (NE Greece)', *Acta Oecologica* 33(1), 27 - 35.
- (392) Vaz, A.; Kerbauy, G. & Figueiredo-Ribeiro, R. (1998), 'Changes in soluble carbohydrates and starch partitioning during vegetative bud formation from root tips of *Catasetum fimbriatum* (Orchidaceae)', *PLANT CELL TISSUE AND ORGAN CULTURE* 54(2), 105-111.
- (393) Vos, P. D. (2010), 'European materia medica in historical texts: Longevity of a tradition and implications for future use', *Journal of Ethnopharmacology* 132(1), 28 - 47.
- (394) Weber, R. W. & Webster, J. (2001), 'Teaching techniques for mycology: 14. Mycorrhizal infection of orchid seedlings in the laboratory', *Mycologist* 15(2), 55 - 59.
- (395) Wells, T. C. E.; Rothery, P.; Cox, R. & Bamford, S. (1998), 'Flowering dynamics of

- Orchis morio L. and Herminium monorchis (L.) R. Br. at two sites in eastern England', Botanical Journal of the Linnean Society 126(1-2), 39 - 48.
- (396) Wells, T.; Rothery, P.; Cox, R. & Bamford, S. (1998), 'Flowering dynamics of Orchis morio L. and Herminium monorchis (L.) R.Br. at two sites in eastern England', BOTANICAL JOURNAL OF THE LINNEAN SOCIETY 126(1-2), 39-48.
- (397) Whipps, J.; Haselwandter, K.; McGee, E. & Lewis, D. (1982), 'Use of biochemical markers to determine growth, development and biomass of fungi in infected tissues, with particular reference to antagonistic and mutualistic biotrophs', Transactions of the British Mycological Society 79(3), 385 - 400.
- (398) Wood, C.; Pritchard, H. & Miller, A. (2000), 'Simultaneous preservation of orchid seed and its fungal symbiont using encapsulation-dehydration is dependent on moisture content and storage temperature', CRYO-LETTERS 21(2), 125-136.
- (399) Wotavová, K.; Balounová, Z. & Kindlmann, P. (2004), 'Factors affecting persistence of terrestrial orchids in wet meadows and implications for their conservation in a changing agricultural landscape', Biological Conservation 118(3), 271 - 279.

ORCHIS USTULATA

- (400) Harastova-Sobotkova, M.; Jersakova, J.; Kindlmann, P. & Curn, L. (2005), 'Morphometric and genetic divergence among populations of *Neotinea ustulata* (Orchidaceae) with different flowering phenologies', *FOLIA GEOBOTANICA* 40(4), 385-405.
- (401) Shefferson, R. P. & Tali, K. (2007), 'Dormancy is associated with decreased adult survival in the burnt orchid, *Neotinea ustulata*', *JOURNAL OF ECOLOGY* 95(1), 217-225.
- (402) Tali, K.; Fay, M. & Bateman, R. (2006), 'Little genetic differentiation across Europe between early-flowering and late-flowering populations of the rapidly declining orchid *Neotinea ustulata*', *BIOLOGICAL JOURNAL OF THE LINNEAN SOCIETY* 87(1), 13-25.
- (403) Tali, K.; Foley, M. J. Y. & Kull, T. (2004), '*Orchis ustulata* L.', *Journal of Ecology* 92(1), 174--184.
- (404) Tali, K. & Kull, T. (2001), 'Highly variable flowering time in *Orchis ustulata* (Orchidaceae): consequences for population dynamics', *NORDIC JOURNAL OF BOTANY* 21(5), 457-466.

PEDICULARIS PALUSTRIS

- (405) Fink, S. (1996), 'Verbreitung, Soziologie und Ökologie von *Pedicularis palustris* L. in Nordwestdeutschland', Master's thesis, Universität Bremen.
- (406) Karrenberg, S. & Jensen, K. (2000), 'Effects of pollination and pollen source on the seed set of *Pedicularis palustris*', *Folia Geobotanica* 35(2), 191-202.
- (407) Koeman-Kwak, M. (1973), 'Pollination of *Pedicularis-palustris* By Nectar Thieves (short-tongued Bumblebees)', *Acta Botanica Neerlandica* 22(6), 608-615.
- (408) Macior, L. W. (1993), 'Pollination ecology of *Pedicularis palustris* L. (Scrophulariaceae) in North America', *Plant Species Biology* 8(1), 35-44.
- (409) Rosenthal, G. & Fink, S. (1996), 'Pedicularis palustris L. in the Bremen area: Distribution, ecology, and causes of its decline', *Abhandlungen Naturwissenschaftlichen Verein zu Bremen* 43(2), 429-447.
- (410) Schmidt, K. & Jensen, K. (2000), 'Genetic structure and AFLP variation of remnant populations in the rare plant *Pedicularis palustris* (Scrophulariaceae) and its relation to population size and reproductive components', *American Journal of Botany* 87(5), 678-689.
- (411) Steffen, K. (1956), 'Endomitosen Im Endosperm Von *Pedicularis Palustris* L', *Planta* 47(6), 625-652.
- (412) Steffen, K. (1955), 'Kern und Nucleolenwachstum bei endomitotischer Polyploidisierung - Ein Beitrag zur karyologischen Anatomie von *Pedicularis palustris* L.', *Planta* 45(4), 379-394.
- (413) Ter Borg, S. J.; Janse, A. & Kwak, M. M. (1980), 'Life Cycle Variation In *Pedicularis palustris* (Scrophulariaceae)', *Acta Botanica Neerlandica* 29(5-6), 397-406.
- (414) Wattez, J. R. (1974), 'Marsh Lousewort *Pedicularis palustris* ssp. *serotina* L In the North of France', *Naturalistes Belges* 55(6), 241-257.

PHYTEUMA ORBICULARE

- (415) Borsch, T.; Korotkova, N.; Raus, T.; Lobin, W. & Loehne, C. (2009), 'The petD group II intron as a species level marker: utility for tree inference and species identification in the diverse genus *Campanula* (Campanulaceae)', *Willdenowia* 39(1), 7--33.
- (416) Dutoit, T.; Buisson, E.; Roche, P. & Alard, D. (2004), 'Land use history and botanical changes in the calcareous hillsides of Upper-Normandy (north-western France): new implications for their conservation management', *Biological Conservation* 115(1), 1 - 19.
- (417) Fischer, M. & Wipf, S. (2002), 'Effect of low-intensity grazing on the species-rich vegetation of traditionally mown subalpine meadows', *Biological Conservation* 104(1), 1 - 11.
- (418) Hurst, A. & John, E. (1999), 'The biotic and abiotic changes associated with *Brachypodium pinnatum* dominance in chalk grassland in south-east England', *Biological Conservation* 88(1), 75 - 84.
- (419) Lassueur, T.; Joost, S. & Randin, C. F. (2006), 'Very high resolution digital elevation models: Do they improve models of plant species distribution?', *Ecological Modelling* 198(1-2), 139 - 153.
- (420) Quinn, R.; Lawton, J.; Eversham, B. & Wood, S. (1994), 'The biogeography of scarce vascular plants in Britain with respect to habitat preference, dispersal ability and reproductive biology', *Biological Conservation* 70(2), 149 - 157.
- (421) Thompson, K. & Hodgson, J. G. (1996), 'More on the biogeography of scarce vascular plants', *Biological Conservation* 75(3), 299 - 302.
- (422) Yu, F.-H.; Krüsi, B.; Schütz, M.; Schneller, J. J. & Wildi, O. (2008), 'Plant communities affect the species-area relationship on *Carex sempervirens* tussocks', *Flora - Morphology, Distribution, Functional Ecology of Plants* 203(3), 197 - 203.

POLYGALA SERPYLLIFOLIA

- (423) Bruelheide, H. (2000), 'Population dynamics of endangered species in a transplanted montane meadow', *Folia Geobotanica* 35(2), 179-189.
- (424) Heubl, G. R. (1984), 'Systematische Untersuchungen an mitteleuropäischen Polygala-Arten. [Systematic Studies of Central European Polygala Species]', *Mitteilungen der Botanischen Staatssammlung Muenchen* 20, 205-428.
- (425) Lack, A. J. (1995), 'Relationships and Hybridization Between British Species of Polygala - Evidence From Isozymes', *New Phytologist* 130(2), 217-223.
- (426) Lack, A. J. & Kay, Q. O. N. (1987), 'Genetic-structure, Gene Flow and Reproductive Ecology In Sand-dune Populations of Polygala vulgaris', *Journal of Ecology* 75(1), 259-276.
- (427) Oostermeijer, J. G. B. (1989), 'Myrmecochory In Polygala-vulgaris L, Luzula-campestris (L) Dc and Viola-curtisii Forster In A Dutch Dune Area', *Oecologia* 78(3), 302-311.
- (428) Trueman, I. C. (1974), 'Variation and Hybridization In British Species of Polygala', *Biological Journal of the Linnean Society* 6(4), 370.

PSEUDORCHIS ALBIDA

- (429) Bateman, R.; Hollingsworth, P.; Preston, J.; Yi-Bo, L.; Pridgeon, A. & Chase, M. (2003), 'Molecular phylogenetics and evolution of Orchidinae and selected Habenariinae (Orchidaceae)', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* 142(1), 1-40.
- (430) Bateman, R.; Rudall, P. & James, K. (2006), 'Phylogenetic context, generic affinities and evolutionary origin of the enigmatic Balkan orchid *Gymnadenia frivaldii* Hampe ex Griseb.', *TAXON* 55(1), 107-118.
- (431) Duffy, K. J.; Kingston, N. E.; Sayers, B. A.; Roberts, D. L. & Stout, J. C. (2009), 'Inferring National and Regional Declines of Rare Orchid Species with Probabilistic Models', *CONSERVATION BIOLOGY* 23(1), 184-195.
- (432) Klein, E. (2000), 'Pseudorchis albida subsp. *tricuspis* (BECK) KLEIN stat. nov., a largely overlooked calcicolous, alpic-boreal taxon (Orchidaceae-Orchideae)', *PHYTON-ANNALES REI BOTANICAE* 40(1), 141-159.
- (433) Kull, T. & Hutchings, M. J. (2006), 'A comparative analysis of decline in the distribution ranges of orchid species in Estonia and the United Kingdom', *Biological Conservation* 129(1), 31 - 39.
- (434) Nantel, P.; Bouchard, A.; Brouillet, L. & Hay, S. (1998), 'Selection of areas for protecting rare plants with integration of land use conflicts: A case study for the west coast of Newfoundland, Canada', *Biological Conservation* 84(3), 223 - 234.
- (435) Rae, D.; Armbruster, W.; Edwards, M. & Svengård-Barre, M. (2006), 'Influence of microclimate and species interactions on the composition of plant and invertebrate communities in alpine northern Norway', *Acta Oecologica* 29(3), 266 - 282.
- (436) Reinhammar, L. & Hedren, M. (1998), 'Allozyme differentiation between lowland and alpine populations of *Pseudorchis albida* s.lat. (Orchidaceae) in Sweden', *NORDIC JOURNAL OF BOTANY* 18(1), 7-14.
- (437) Reinhammar, L. (1995), 'Evidence for two distinctive species of *Pseudorchis* (Orchidaceae) in Scandinavia.', *NORDIC JOURNAL OF BOTANY* 15(5), 469-481.
- (438) Reinhammar, L. (1998), 'Systematics of *Pseudorchis albida* sl (Orchidaceae) in Europe and North America', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* 126(4), 363-382.
- (439) Reinhammar, L.; Olsson, E. & Sormeland, E. (2002), 'Conservation biology of an endangered grassland plant species, *Pseudorchis albida*, with some references to the closely related alpine *P-straminea* (Orchidaceae)', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* 139(1), 47-66.
- (440) Reinhammar, L.-G. & Hedren, M. (1998), 'Allozyme differentiation between lowland and alpine populations of *Pseudorchis albida* s.lat. (Orchidaceae) in Sweden', *Nordic*

Journal of Botany 18(1), 7--14.

- (441) Reinhammar, L.-G.; Olsson, E. G. A. & Sormeland, E. (2002), 'Conservation biology of an endangered grassland plant species, *Pseudorchis albida*, with some references to the closely related alpine *P. straminea* (Orchidaceae)', *Botanical Journal of the Linnean Society* 139, 47-66.
- (442) SCHWABE, A. (1990), 'Syndynamic Processes In Nardo-callunetea Communities Changes In Fallow Land After Renewed Cattle Grazing and Life History of *Arnica montana* L', *Carolinea* 48, 45--68.
- (443) Yeo, M. J. M.; Blackstock, T. H. & Stevens, D. P. (1998), 'The use of phytosociological data in conservation assessment: a case study of lowland grasslands in mid Wales', *Biological Conservation* 86(2), 125 - 138.

SWERTIA PERENNIS

- (444) Kirichok, E. I. (1999), 'The record of *Swertia perennis* (Gentianaceae) northwest of Smolensk region', *Botanicheskii Zhurnal* (St. Petersburg) 84(2), 124-127.
- (445) Lienert, J.; Diemer, M. & Schmid, B. (2002), 'Effects of habitat fragmentation on population structure and fitness components of the wetland specialist *Swertia perennis* L. (Gentianaceae)', *Basic and Applied Ecology* 3(2), 101-114.
- (446) Lienert, J. & Fischer, M. (2004), 'Experimental inbreeding reduces seed production and germination independent of fragmentation of populations of *Swertia perennis*', *Basic and Applied Ecology* 5(1), 43-52.
- (447) Lienert, J.; Fischer, M. & Diemer, M. (2002), 'Local extinctions of the wetland specialist *Swertia perennis* L. (Gentianaceae) in Switzerland: A revisitation study based on herbarium records', *Biological Conservation* 103(1), 65-76.
- (448) Lienert, J.; Fischer, M.; Schneller, J. & Diemer, M. (2002), 'Isozyme variability of the wetland specialist *Swertia perennis* (Gentianaceae) in relation to habitat size, isolation, and plant fitness', *American Journal of Botany* 89(5), 801-811.
- (449) Romero, M. T. & Rico, H. E. (1986), '*Swertia-perennis* New-record L. For the Central Mountain System Spain', *Anales del Jardin Botanico de Madrid* 43(1), 177-178.
- (450) Stoicovici, L. (1984), 'Interspecific Relationships of the Glacial Relics *Swertia-perennis* L and *Pedicularis-sceptrum-carolinum* L In A Romanian Fen', *Vegetatio* 56(3), 139-145.

TRIFOLIUM MONTANUM

- (451) Bulinska-Radomska, Z. (2000), 'Morphological relationships among 15 species of Trifolium occurring in Poland', *Genetic Resources and Crop Evolution* 47(3), 267--272.
- (452) Court-Picon, M.; Buttler, A. & de Beaulieu, J.-L. (2005), 'Modern pollen-vegetation relationships in the Champsaur valley (French Alps) and their potential in the interpretation of fossil pollen records of past cultural landscapes', *Review of Palaeobotany and Palynology* 135(1-2), 13 - 39.
- (453) Cousins, S. A. & Aggemyr, E. (2008), 'The influence of field shape, area and surrounding landscape on plant species richness in grazed ex-fields', *Biological Conservation* 141(1), 126 - 135.
- (454) Dear, B. S.; Reed, K. F. M. & Craig, A. D. (2008), 'Outcomes of the search for new perennial and salt tolerant pasture plants for southern Australia', *Australian Journal of Experimental Agriculture* 48(4), 578--588.
- (455) Gostin, I. N. (2009), 'Air Pollution Effects on the Leaf Structure of some Fabaceae Species', *Notulae Botanicae Horti Agrobotanici Cluj- Napoca* 37(2), 57--63.
- (456) Gucwa-Przepiora, E. & Blaszkowski, J. (2007), 'Arbuscular mycorrhiza of plants spontaneously colonizing the soda heap in Jaworzno (Southern Poland)', *Acta Societatis Botanicorum Poloniae* 76(1), 69--74.
- (457) Herzog, F.; Dreier, S.; Hofer, G.; Marfurt, C.; Schüpbach, B.; Spiess, M. & Walter, T. (2005), 'Effect of ecological compensation areas on floristic and breeding bird diversity in Swiss agricultural landscapes', *Agriculture, Ecosystems & Environment* 108(3), 189 - 204.
- (458) KAZAKOV, A. Z. (1977), 'Flavonoids of Trifolium-montanum', *Khimiya Prirodnikh Soedinenii*(3), 415 - 416.
- (459) KAZIMIERSKA, E. M. & KAZIMIERSKI, T. (1985), 'Crossing of Species In the Genus Trifolium L. 1. Trifolium-repens Lxt-montanum L - Fertilization and Development of the Embryo and Endosperm', *Genetica Polonica* 26(3), 327--.
- (460) Klimes[breve]ová, J. & Klimes, L. (2007), 'Bud banks and their role in vegetative regeneration - A literature review and proposal for simple classification and assessment', *Perspectives in Plant Ecology, Evolution and Systematics* 8(3), 115 - 129.
- (461) Lauber, W. & Körner, C. (1997), 'In situ stomatal responses to long-term CO₂ enrichment in calcareous grassland plants', *Acta Oecologica* 18(3), 221 - 229.
- (462) Marini, L.; Scotton, M.; Klimek, S.; Isselstein, J. & Pecile, A. (2007), 'Effects of local factors on plant species richness and composition of Alpine meadows', *Agriculture, Ecosystems & Environment* 119(3-4), 281 - 288.

- (463) Menzano, A.; Rambozzi, L. & Rossi, L. (2007), 'A severe episode of wildlife-derived scabies in domestic goats in Italy', *Small Ruminant Research* 70(2-3), 154 - 158.
- (464) Muncaciu, S.; Gafta, D.; Cristea, V.; Rosca-Casian, O. & Goia, I. (2010), 'Eco-coenotic conditions and structure of *Trollius europaeus* L. populations in an extrazonal habitat complex (Transylvanian Carpathian foothills)', *Flora - Morphology, Distribution, Functional Ecology of Plants* In Press, Corrected Proof, - .
- (465) Oklejewicz, K. (1997), 'Veronica gentianoides (Scrophulariaceae): An interesting synanthropic species in the Polish flora', *Fragmenta Floristica et Geobotanica Series Polonica* 4(0), 377--378.
- (466) Piqueray, J.; Bottin, G.; Delescaille, L.-M.; Bisteau, E.; Colinet, G. & Mahy, G. (2010), 'Rapid restoration of a species-rich ecosystem assessed from soil and vegetation indicators: The case of calcareous grasslands restored from forest stands', *Ecological Indicators* In Press, Corrected Proof, - .
- (467) Przywara, L.; Pirog, H.; Konieczny, R. & Jach, M. (1996), 'The use of tissue culture methods in interspecific hybridization within the genus *Trifolium* L.', *Acta Biologica Cracoviensia Series Botanica* 38, 53--.
- (468) Reiné, R.; Chocarro, C. & Fillat, F. (2004), 'Soil seed bank and management regimes of semi-natural mountain meadow communities', *Agriculture, Ecosystems & Environment* 104(3), 567 - 575.
- (469) Schleuning, M. & Matthies, D. (2009), 'Habitat Change and Plant Demography: Assessing the Extinction Risk of a Formerly Common Grassland Perennial', *Conservation Biology* 23(1), 174--183.
- (470) Schleuning, M.; Niggemann, M.; Becker, U. & Matthies, D. (2009), 'Negative effects of habitat degradation and fragmentation on the declining grassland plant *Trifolium montanum*', *Basic and Applied Ecology* 10(1), 61 - 69.
- (471) Schwabe, A.; Kratochwil, A. & Pignatti, S. (2007), 'Plant indicator values of a high-phytodiversity country (Italy) and their evidence, exemplified for model areas with climatic gradients in the southern inner Alps', *Flora - Morphology, Distribution, Functional Ecology of Plants* 202(5), 339 - 349.
- (472) Spitzer, L.; Benes, J.; Dandova, J.; Jaskova, V. & Konvicka, M. (2009), 'The Large Blue butterfly, *Phengaris* [Maculinea] *arion*, as a conservation umbrella on a landscape scale: The case of the Czech Carpathians', *Ecological Indicators* 9(6), 1056 - 1063.
- (473) Sudova, R. (2009), 'Different growth response of five co-existing stoloniferous plant species to inoculation with native arbuscular mycorrhizal fungi', *Plant Ecology* 204(1), 135--143.
- (474) van der Merwe, M. M.; Walker, J.; Ericson, L. & Burdon, J. J. (2008), 'Coevolution with higher taxonomic host groups within the *Puccinia/Uromyces* rust lineage obscured by

- host jumps', *Mycological Research* 112(12), 1387 - 1408.
- (475) van der Merwe, M.; Ericson, L.; Walker, J.; Thrall, P. H. & Burdon, J. J. (2007), 'Evolutionary relationships among species of *Puccinia* and *Uromyces* (Pucciniaceae, Uredinales) inferred from partial protein coding gene phylogenies', *Mycological Research* 111(2), 163 - 175.
- (476) VETTER, J. (1995), 'Isoflavones In Different Parts of Common *Trifolium* Species', *Journal of Agricultural and Food Chemistry* 43(1), 106--108.
- (477) Vilcinskas, E. & Dabkeviciene, G. (2009), 'Qualitative and quantitative characteristics of clover (*Trifolium* spp.) species in the first year of growing', *Zemdirbyste-agriculture* 96(4), 170--180.
- (478) Wohlfahrt, G.; Bahn, M.; Horak, I.; Tappeiner, U. & Cernusca, A. (1998), 'A nitrogen sensitive model of leaf carbon dioxide and water vapour gas exchange: application to 13 key species from differently managed mountain grassland ecosystems', *Ecological Modelling* 113(1-3), 179 - 199.

GESAMTÜBERSICHT

- (1) Abadie, J. C.; Puttsepp, U.; Gebauer, G.; Faccio, A.; Bonfante, P. & Selosse, M. A. (2006), 'Cephalanthera longifolia (Neottieae, Orchidaceae) is mixotrophic: a comparative study between green and nonphotosynthetic individuals', *Canadian Journal of Botany-revue Canadienne De Botanique* **84**(9), 1462--1477.
- (2) Aceto, S.; Cozzolino, S.; Gaudio, L.; Nazzaro, R. & De Luca, P. (1999), 'Pollination flow in hybrid formation between *Orchis morio* and *Orchis papilionacea* (Orchidaceae) in two different habitats', *INTERNATIONAL JOURNAL OF PLANT SCIENCES* **160**(6), 1153-1156.
- (3) Aceto, S.; Montieri, S.; Sica, M. & Gaudio, L. (2007), 'Molecular evolution of the OrcPI locus in natural populations of Mediterranean orchids', *Gene* **392**(1-2), 299 - 305.
- (4) Adriaens, D.; Honnay, O. & Hermy, M. (2006), 'No evidence of a plant extinction debt in highly fragmented calcareous grasslands in Belgium', *Biological Conservation* **133**(2), 212 - 224.
- (5) Ahmed, A. A.; Hussein, T. A.; Mahmoud, A. A.; Farag, M. A.; Pare, P. W.; Wojcinska, M.; Karchesy, J. & Mabry, T. J. (2004), 'Nor-ent-kaurane diterpenes and hydroxylactones from *Antennaria geyeri* and *Anaphalis margaritacea*', *Phytochemistry* **65**(18), 2539--2543.
- (6) Aksenova, A. A.; Onipchenko, V. G. & Blinnikov, M. S. (1998), 'Plant interactions in alpine tundra: 13 years of experimental removal of dominant species', *Ecoscience* **5**(2), 258--270.
- (7) Ale-Agha, N.; Feige, G. B.; Jensen, M.; Christiaans, B.; Brassmann, M. & Kricke, R. (2004), 'New, rare or remarkable microfungi in the Italian Alps (Carnic Alps) - Part II - Other microfungi', *Communications in Agricultural and Applied Biological Sciences* **69**(4), 467--477.
- (8) Alonso-Sande, M.; Teijeiro-Osorio, D.; Remuñán-López, C. & Alonso, M. (2009), 'Glucomannan, a promising polysaccharide for biopharmaceutical purposes', *European Journal of Pharmaceutics and Biopharmaceutics* **72**(2), 453 - 462.
- (9) Alpaslan, M. & Hayta, M. (2007), 'Effect of soymilk substitution on the rheological and sensory properties of salep (Traditional turkish milk beverage)', *INTERNATIONAL JOURNAL OF FOOD PROPERTIES* **10**(3), 413-420.
- (10) Andersson, U.-B. (2004), 'Botany day in Orebo county 2004', *Svensk Botanisk Tidskrift* **98**(5), 263--267.
- (11) Atalla, R.; Hackney, J.; Uhlin, I. & Thompson, N. (1993), 'Hemicelluloses as structure regulators in the aggregation of native cellulose', *International Journal of Biological Macromolecules* **15**(2), 109 - 112.
- (12) Austrheim, G.; Gunilla, E.; Olsson, A. & Grøntvedt, E. (1999), 'Land-use impact on plant communities in semi-natural sub-alpine grasslands of Budalen, central Norway', *Biological Conservation* **87**(3), 369 - 379.

- (13) Aybeke, M.; Sezik, E. & Olgun, G. (2010), 'Vegetative anatomy of some Ophrys, Orchis and Dactylorhiza (Orchidaceae) taxa in Trakya region of Turkey', *Flora - Morphology, Distribution, Functional Ecology of Plants* **205**(2), 73 - 89.
- (14) Aziz, N.; Mehmood, M. H.; Siddiqi, H. S.; Mandukhail, S.-u.-R.; Sadiq, F.; Maan, W. & Gilani, A. H. (2009), 'Antihypertensive, antidiabetic and endothelial modulating activities of *Orchis mascula*', *HYPERTENSION RESEARCH* **32**(11), 997-1003.
- (15) Badria, F. A. (1994), 'Is man helpless against cancer? An environmental approach: antimutagenic agents from Egyptian food and medicinal preparations', *Cancer Letters* **84**(1), 1 - 5.
- (16) Baltisberger, M. & Widmer, A. (2009), 'Karyological data of some angiosperms from Romania', *Willdenowia* **39**(2), 353--363.
- (17) Barbaro, L.; Dutoit, T.; Anthelme, F. & Corcket, E. (2004), 'Respective influence of habitat conditions and management regimes on prealpine calcareous grasslands', *Journal of Environmental Management* **72**(4), 261 - 275.
- (18) Barkham, J. (1992), 'The effects of management on the ground flora of ancient woodland, Brigsteer Park Wood, Cumbria, England', *Biological Conservation* **60**(3), 167 - 187.
- (19) Barreto, D. W. & Parente, J. P. (2006), 'Chemical properties and biological activity of a polysaccharide from *Cyrtopodium cardiochilum*', *Carbohydrate Polymers* **64**(2), 287 - 291.
- (20) Bartoszek, W. (1999), 'A new station for *Carex pulicaris* (Cyperaceae) in the Polish Carpathians (Nowe stanowisko *Carex pulicaris* (Cyperaceae) w Karpatach polskich)', *Fragmenta Floristica et Geobotanica - Series Polonica* **6**(0), 274--277.
- (21) Baskin, C. C. & Baskin, J. M. (1998), *Germination Ecology of Plants with Specialized Life Cycles and/or Habitats*'Seeds', Academic Press, San Diego, pp. 459 - 557.
- (22) Bateman, R.; Hollingsworth, P.; Preston, J.; Yi-Bo, L.; Pridgeon, A. & Chase, M. (2003), 'Molecular phylogenetics and evolution of Orchidinae and selected Habenariinae (Orchidaceae)', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* **142**(1), 1-40.
- (23) Bateman, R.; Rudall, P. & James, K. (2006), 'Phylogenetic context, generic affinities and evolutionary origin of the enigmatic Balkan orchid *Gymnadenia frivaldii* Hampe ex Griseb.', *TAXON* **55**(1), 107-118.
- (24) Bayer, R. J. (1993), 'A Taxonomic Revision of the Genus *Antennaria* (Asteraceae: Inuleae: Gnaphaliinae) of Alaska and Yukon Territory, Northwestern North America', *Arctic and Alpine Research* **25**(2), 150-159.
- (25) Bayer, R. J. (1989), 'A Taxonomic Revision of the *Antennaria rosea* (Asteraceae: Inuleae: Gnaphaliinae) Polyploid Complex', *Brittonia* **41**(1), 53-60.

- (26) Bayer, R. J. (1984), 'Chromosome Numbers and Taxonomic Notes for North American Species of *Antennaria* (Asteraceae: Inuleae)', *Systematic Botany* **9**(1), 74-83.
- (27) Bayer, R. J.; Soltis, D. E. & Soltis, P. S. (1996), 'Phylogenetic Inferences in *Antennaria* (Asteraceae: Gnaphalieae: Cassiniinae) Based on Sequences from Nuclear Ribosomal DNA Internal Transcribed Spacers (ITS)', *American Journal of Botany* **83**(4), 516-527.
- (28) Bayer, R. J. & Stebbins, G. L. (1987), 'Chromosome Numbers, Patterns of Distribution, and Apomixis in *Antennaria* (Asteraceae: Inuleae)', *Systematic Botany* **12**(2), 305-319.
- (29) Becker, U.; Colling, G.; Dostal, P.; Jakobsson, A. & Matthies, D. (2006), 'Local adaptation in the monocarpic perennial *Carlina vulgaris* at different spatial scales across Europe', *Oecologia* **150**(3), 506-518.
- (30) Benincasa, M.-A.; Cartoni, G. & Fratte, C. D. (2002), 'Flow field-flow fractionation and characterization of ionic and neutral polysaccharides of vegetable and microbial origin', *Journal of Chromatography A* **967**(2), 219 - 234.
- (31) van den Berg, L. J. L.; Peters, C. J. H.; Ashmore, M. R. & Roelofs, J. G. M. (2008), 'Reduced nitrogen has a greater effect than oxidised nitrogen on dry heathland vegetation', *Environmental Pollution* **154**(3), 359--369.
- (32) Bergamini, A.; Peintinger, M.; Fakheran, S.; Moradi, H.; Schmid, B. & Joshi, J. (2009), 'Loss of habitat specialists despite conservation management in fen remnants 1995-2006', *Perspectives in Plant Ecology, Evolution and Systematics* **11**(1), 65 - 79.
- (33) Berlin, G. A. I.; Linusson, A.-C. & Olsson, E. G. A. (2000), 'Vegetation changes in semi-natural meadows with unchanged management in southern Sweden, 1965-1990', *Acta Oecologica* **21**(2), 125 - 138.
- (34) BEYRLER, H.; SMITH, S.; PETERSON, R. & FRANCO, C. (1995), 'COLONIZATION OF ORCHIS-MORIO PROTOCOLS BY A MYCORRHIZAL FUNGUS - EFFECTS OF NITROGEN NUTRITION AND GLYPHOSATE IN MODIFYING THE RESPONSES', *CANADIAN JOURNAL OF BOTANY-REVUE CANADIENNE DE BOTANIQUE* **73**(8), 1128-1140.
- (35) BICCHI, C.; NANO, G. M. & TIRA, S. (1975), 'Normal-paraffin Components of Some Gnaphalieae', *Planta Medica* **28**(4), 389--391.
- (36) Blachnik, T. (2010), 'Artenhilfsprojekt Arnika und Katzenpfötchen im Bayerischen Vogtland, Landkreis Hof - Fachgrundlagen - Verbreitung, Bestandssituation, Erhaltungsmaßnahmen und Öffentlichkeitsarbeit für Leitarten bodensaurer Magerrasen', Technical report, Agentur und Naturschutzbüro Blachnik.
- (37) Blachnik, T. (2009), 'Arnika und Katzenpfötchen - Ein Artenhilfsprojekt im Bayerischen Vogtland', Herausgeber:Regierung von Oberfranken.
- (38) Blackmore, S. (1984), 'Compositae - lactuceae', *Review of Palaeobotany and Palynology* **42**(1-

- 4), 45 - 85.
- (39) Blinnikov, M. S. (1994), 'Experimental investigation of alpine plant communities in the Northwestern Caucasus: Phytolith analysis and holocene dynamics of alpine vegetation', *Veroeffentlichungen des Geobotanischen Institutes der Eidgenoessische Technische Hochschule Stiftung Ruebel in Zuerich* **0**(115), 23--40.
- (40) Bobbink, R. & Roelofs, J. G. (1995), 'Ecological effects of atmospheric deposition on non-forest ecosystems in Western Europe', *Studies in Environmental Science* **64**, 279 - 292.
- (41) Boehling, N. (2003), 'Investigations of permanent plots in an oak-hornbeam woodland in the foothills of the Swabian mountains (SW Germany), 1978-2001: Decline of *Scilla bifolia* and the invasion of *Allium ursinum*.', *Tuexenia*(23), 131--161.
- (42) Bohlin, K. (2006), 'Different colour forms in *Antennaria dioica*', *Svensk Botanisk Tidskrift* **100**(1), 3--4.
- (43) Bohlmann, F. & Zdero, C. (1971), 'Terpene derivatives from higher plants .11. prutenin, 4-angeloyloxypruteninone, and 4-acetoxypruteninone and 4-acetoxyisopruteninone - 4 new sesquiterpene lactons from *Laserpitium-prutenicum* l', *chemische Berichte-recueil* **104**(5), 1611-.
- (44) Borsch, T.; Korotkova, N.; Raus, T.; Lobin, W. & Loehne, C. (2009), 'The petD group II intron as a species level marker: utility for tree inference and species identification in the diverse genus *Campanula* (Campanulaceae)', *Willdenowia* **39**(1), 7--33.
- (45) Braeuer, I.; Maibom, W.; Matthies, D.; Tschardt, T. & fuer Oekologie, G. (1998), 'Population size and risk of extinction in endangered plant species in Lower Saxony', *Verhandlungen der Gesellschaft fuer Oekologie*, 505--510.
- (46) Breadmore, K. N. & Kirk, W. D. J. (1998), 'Factors affecting floral herbivory in a limestone grassland', *Acta Oecologica* **19**(6), 501 - 506.
- (47) BROWN, L. S. R. & GRAY, D. O. (1988), '1-amino-2-propanol, A Natural Product From the Compositae', *Phytochemistry* **27**(4), 1195--1197.
- (48) Buchala, A. J.; Franz, G. & Meier, H. (1974), 'A glucomannan from the tubers of *Orchis morio*', *Phytochemistry* **13**(1), 163 - 166.
- (49) BUHL, O.; FALCK, P.; KARSHOLT, O.; LARSEN, K. & SCHNACK, K. (1989), 'Records of Microlepidoptera From Denmark In 1987 Lepidoptera', *Entomologische Meddelelser* **57**(3), 123--135.
- (50) Bulinska-Radomska, Z. (2000), 'Morphological relationships among 15 species of *Trifolium* occurring in Poland', *Genetic Resources and Crop Evolution* **47**(3), 267--272.
- (51) Bullock, J. M. (1998), 'Community translocation in Britain: Setting objectives and measuring

- consequences', *Biological Conservation* **84**(3), 199 - 214.
- (52) Bundesregierung (), 'BNatSchG'.
- (53) Böhnert, W. (1991), 'Die Orchidee des Jahres 1991. Das Kleine Knabenkraut (*Orchis morio*).', *Ber. Arbeitskrs. Heim. Orchid.* **8**(1), 86-91.
- (54) Böhnert, W. & Hamel, G. (1988), 'Zur gegenwärtigen Situation des Kleinen Knabenkrautes (*Orchis morio* L.) in der DDR - Populationsituation, Schutz und Betreuung', *Arch. Naturschutz und Landschafts.Forsch.* **28**(2), 101-119.
- (55) Cafasso, D.; Pellegrino, G.; Caputo, P.; Scrugli, A. & Cozzolino, S. (2001), 'Genetic relatedness of insular segregates of mediterranean orchid species as inferred from ITS sequences analysis', *BOTANICA HELVETICA* **111**(2), 181-190.
- (56) Calster, H. V.; Endels, P.; Antonio, K.; Verheyen, K. & Hermy, M. (2008), 'Coppice management effects on experimentally established populations of three herbaceous layer woodland species', *Biological Conservation* **141**(10), 2641 - 2652.
- (57) Cantone, C.; Sica, M.; Gaudio, L. & Aceto, S. (2009), 'The *OrcPI* locus: Genomic organization, expression pattern, and noncoding regions variability in *Orchis italica* (Orchidaceae) and related species', *Gene* **434**(1-2), 9 - 15.
- (58) Caputo, P.; Aceto, S.; Cozzolino, S. & Nazzaro, R. (1997), 'Morphological and molecular characterization of a natural hybrid between *Orchis laxiflora* and *O-morio* (Orchidaceae)', *PLANT SYSTEMATICS AND EVOLUTION* **205**(3-4), 147-155.
- (59) Carruthers, W.; Duncan, F. & Shrubsole, W. (1875), 'Excursion to the isle of Sheppey: June 7th, 1875', *Proceedings of the Geologists' Association* **4**(5), 320 - 323.
- (60) Cescutti, P.; Campa, C.; Delben, F. & Rizzo, R. (2002), 'Structure of the oligomers obtained by enzymatic hydrolysis of the glucomannan produced by the plant *Amorphophallus konjac*', *Carbohydrate Research* **337**(24), 2505 - 2511.
- (61) Cescutti, P.; Pupulin, R.; Delben, F.; Abbate, M.; Dentini, M.; Sparapano, L.; Rizzo, R. & Crescenzi, V. (2002), 'New exopolysaccharides produced by *Aureobasidium pullulans* grown on glucosamine', *Carbohydrate Research* **337**(13), 1203 - 1209.
- (62) Challe, J. & Struik, P. (2008), 'The impact on orchid species abundance of gathering their edible tubers by HIV/AIDS orphans: a case of three villages in the Southern Highlands of Tanzania', *NJAS - Wageningen Journal of Life Sciences* **56**(3), 261 - 279.
- (63) Chiryat'ev, E. A. & Rusakova, O. A. (1994), 'Species of Siberian flora as sources of direct action anti-coagulation', *Rastitel'nye Resursy* **30**(4), 21--28.
- (64) Chmielewski, J. G. (1998), '*Antennaria dioica* (Asteraceae: Inuleae): Addition to the vascular flora of California', *Madrono* **45**(3), 271--272.

- (65) CHMIELEWSKI, J. G.; CHINNAPPA, C. C. & SEMPLE, J. C. (1990), 'The Genus *Antennaria* (asteraceae, Inuleae) In Western North-america - Morphometric Analysis of *Antennaria-alborosea*, *Antennaria-corymbosa*, *Antennaria-marginata*, *Antennaria-microphylla*, *Antennaria-parvifolia*, *Antennaria-rosea*, and *Antennaria-umbrinella*', *Plant Systematics and Evolution* **169**(3-4), 151--175.
- (66) Chrungoo, N. (1992), 'Concepts of dormancy regulation in vegetative plant propagules: A review', *Environmental and Experimental Botany* **32**(4), 309 - 318.
- (67) Coates, F.; Lunt, I. D. & Tremblay, R. L. (2006), 'Effects of disturbance on population dynamics of the threatened orchid *Prasophyllum correctum* D.L. Jones and implications for grassland management in south-eastern Australia', *Biological Conservation* **129**(1), 59 - 69.
- (68) Collin, C. L. & Shykoff, J. A. (2003), 'Outcrossing rates in the gynomonoeocious-gynodioecious species *Dianthus sylvestris* (Caryophyllaceae)', *Am. J. Bot.* **90**(4), 579-585.
- (69) Cooke, D. (2002), 'Possible origin of rampant mignonette populations from garden escapes.', *Plant Protection Quarterly* **17**(4), 147--148.
- (70) Coroi, M.; Skeffington, M. S.; Giller, P.; Smith, C.; Gormally, M. & O'Donovan, G. (2004), 'Vegetation diversity and stand structure in streamside forests in the south of Ireland', *Forest Ecology and Management* **202**(1-3), 39 - 57.
- (71) Court-Picon, M.; Buttler, A. & de Beaulieu, J.-L. (2005), 'Modern pollen-vegetation relationships in the Champsaur valley (French Alps) and their potential in the interpretation of fossil pollen records of past cultural landscapes', *Review of Palaeobotany and Palynology* **135**(1-2), 13 - 39.
- (72) Cousins, S. A. & Aggemyr, E. (2008), 'The influence of field shape, area and surrounding landscape on plant species richness in grazed ex-fields', *Biological Conservation* **141**(1), 126 - 135.
- (73) Cozzolino, S.; Aceto, S.; Caputo, P.; Gaudio, L. & Nazzaro, R. (1998), 'Phylogenetic relationships in *Orchis* and some related genera: an approach using chloroplast DNA', *NORDIC JOURNAL OF BOTANY* **18**(1), 79-87.
- (74) Cozzolino, S.; Nardella, A.; Impagliazzo, S.; Widmer, A. & Lexer, C. (2006), 'Hybridization and conservation of Mediterranean orchids: Should we protect the orchid hybrids or the orchid hybrid zones?', *BIOLOGICAL CONSERVATION* **129**(1), 14-23.
- (75) Cozzolino, S.; Schiestl, F.; Muller, A.; De Castro, O.; Nardella, A. & Widmer, A. (2005), 'Evidence for pollinator sharing in Mediterranean nectar-mimic orchids: absence of pre-mating barriers?', *PROCEEDINGS OF THE ROYAL SOCIETY B-BIOLOGICAL SCIENCES* **272**(1569), 1271-1278.
- (76) Cozzolino, S. & Widmer, A. (2005), 'Orchid diversity: an evolutionary consequence of deception?', *Trends in Ecology & Evolution* **20**(9), 487 - 494.
- (77) Cranston, D. M. & Valentine, D. H. (1983), 'Transplant experiments on rare plant species from

- Upper Teesdale', *Biological Conservation* **26**(2), 175 - 191.
- (78) CRAVEN, A. F. & CRAVEN, S. M. (1977), 'A 2nd Station For *Diapensia-laponica* New-record', *Watsonia* **11**(4), 376.
- (79) D'Emerico, S.; Cozzolino, S.; Pellegrino, G.; Pignone, D. & Scrugli, A. (2002), 'Heterochromatin distribution in selected taxa of the 42-chromosomes *Orchis* s. l. (Orchidaceae)', *CARYOLOGIA* **55**(1), 55-62.
- (80) D'Emerico, S.; Galasso, I.; Pignone, D. & Scrugli, A. (2001), 'Localization of rDNA loci by fluorescent in situ hybridization in some wild orchids from Italy (Orchidaceae)', *CARYOLOGIA* **54**(1), 31-36.
- (81) Darkwa, A. A. & Scott, P. (2009), 'Seasonal growth of temperate terrestrial orchids as affected by the old tuber', *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology* **153**(2, Supplement 1), Abstracts of the Annual Main Meeting of the Society of Experimental Biology, 28th June - 1st July, Glasgow, UK, S212 - S212.
- (82) Darkwa, A. & Scott, P. (2008), 'Growth and influence of defoliation in terrestrial orchids, *Anacamptis morio* and *Serapia lingua*', *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology* **150**(3, Supplement 1), Abstracts of the Annual Main Meeting of the Society of Experimental Biology, 6th - 10th July 2008, Marseille, France, S197 - S197.
- (83) DAVYDOVA, M. V. (1982), 'Effect of Factory Smoke On Grassy Pine Forests In the Southern Urals Russian-sfsr Ussr', *Biologicheskie Nauki (Moscow)*(11), 71--74.
- (84) Dear, B. S.; Reed, K. F. M. & Craig, A. D. (2008), 'Outcomes of the search for new perennial and salt tolerant pasture plants for southern Australia', *Australian Journal of Experimental Agriculture* **48**(4), 578--588.
- (85) DEMERICO, S.; BIANCO, P. & MEDAGLI, P. (1993), 'CYTOLOGICAL AND KARYOLOGICAL STUDIES ON ORCHIDACEAE', *CARYOLOGIA* **46**(4), 309-319.
- (86) D'Emerico, S.; Bianco, P. & Pignone, D. (1996), 'Cytomorphological characterization of diploid and triploid individuals of *Orchis x gennarii* Reichenb fil (Orchidaceae)', *CARYOLOGIA* **49**(2), 153-161.
- (87) D'Emerico, S.; Pignone, D. & Bianco, P. (1996), 'Karyomorphological analyses and heterochromatin characteristics disclose phyletic relationships among 2n=32 and 2n=36 species of *Orchis* (Orchidaceae)', *PLANT SYSTEMATICS AND EVOLUTION* **200**(1-2), 111-124.
- (88) Dersch, G. & Mast, R. (2000), 'Distribution and community formation by *Calamagrostis phragmitoides* HARTMAN (Scandinavian Small-reed, Poaceae) in the German Harz Mountains', *Tuexenia* **20**(20), 119-129.
- (89) Dietrich, W. (2006), 'Contribution to the knowledge of phytopathogenic micromycetes in

- Saxony', *Boletus* **28**(2), 109--118.
- (90) Dietrich, W. (2005), 'The rust fungi, smut fungi and downy mildews in the Czech part of Krusnohory (Erzgebirge): first supplement', *Czech Mycology* **57**(3-4), 257--273.
- (91) Dijk, E. (1990), 'Effects of mycorrhizal fungi on in-vitro nitrogen response of juvenile orchids', *Agriculture, Ecosystems & Environment* **29**(1-4), 91 - 97.
- (92) DIJK, E. & ECK, N. (1995), 'EFFECTS OF MYCORRHIZAL FUNGI ON IN-VITRO NITROGEN RESPONSE OF SOME DUTCH INDIGENOUS ORCHID SPECIES', *CANADIAN JOURNAL OF BOTANY-REVUE CANADIENNE DE BOTANIQUE* **73**(8), 1203-1211.
- (93) Dijk, E. & Grootjans, A. (1998), 'Performance of four *Dactylorhiza* species over a complex trophic gradient', *ACTA BOTANICA NEERLANDICA* **47**(3), 351-368.
- (94) div (2008), 'Expertenworkshop „Biodiversität der Graslandökosysteme Mitteleuropas““Kurzfassungen der Vorträge', Andreas Hermes Akademie Bonn-Rüttgen.
- (95) Dony, J. G. (1977), 'Change in the Flora of Bedfordshire, England, from 1798 to 1976', *Biological Conservation* **11**(4), 307 - 320.
- (96) Dormont, L.; Delle-Vedove, R.; Bessiere, J. M.; Hossaert-Mc Key, M. & Schatz, B. (2010), 'Rare white-flowered morphs increase the reproductive success of common purple morphs in a food-deceptive orchid', *NEW PHYTOLOGIST* **185**(1), 300-310.
- (97) Dragana, S.; Comic, L.; Stefanovic, O. & Solujic-Sukdolac, S. (2008), 'In vitro antibacterial activity of *Antennaria dioica* L. and evaluation of synergism between plant extract and preservatives', *Planta Medica* **74**(9), 1206--1206.
- (98) DUECK, T. A.; VANDEREERDEN, L. J. & BERDOWSKI, J. J. M. (1992), 'Estimation of So₂ Effect Thresholds For Heathland Species', *Functional Ecology* **6**(3), 291--296.
- (99) Duffy, K. J.; Kingston, N. E.; Sayers, B. A.; Roberts, D. L. & Stout, J. C. (2009), 'Inferring National and Regional Declines of Rare Orchid Species with Probabilistic Models', *CONSERVATION BIOLOGY* **23**(1), 184-195.
- (100) Dunnett, N.; Nagase, A.; Booth, R. & Grime, P. (2008), 'Influence of vegetation composition on runoff in two simulated green roof experiments', *Urban Ecosystems* **11**(4), 385--398.
- (101) Dutoit, T.; Buisson, E.; Roche, P. & Alard, D. (2004), 'Land use history and botanical changes in the calcareous hillsides of Upper-Normandy (north-western France): new implications for their conservation management', *Biological Conservation* **115**(1), 1 - 19.
- (102) DVORAK, F. (1976), 'Study of the Number of Chromosomes of 4 Angiosperms', *Scripta Facultatis Scientiarum Naturalium Universitatis Purkynianae Brunensis* **6**(2-3), 113--138.
- (103) Effmert, U.; Saschenbrecker, S.; Ross, J.; Negre, F.; Fraser, C. M.; Noel, J. P.; Dudareva, N. &

- Piechulla, B. (2005), 'Floral benzenoid carboxyl methyltransferases: From in vitro to in planta function', *Phytochemistry* **66**(11), 1211 - 1230.
- (104) EHRENDORFER, F.; LUFTENSTEINER, H. W. & PETRAK, J. (1980), 'Wind Tunnels For the Investigation of Anemochorous Dispersal Units', *Plant Systematics and Evolution* **136**(1-2), 1--6.
- (105) El Naggar, S. M. (2002), 'Taxonomic significance of pollen morphology in some taxa of Resedaceae.', *Feddes Repertorium* **113**(7-8), 518--527.
- (106) Eriksson, O. (1997), 'Colonization dynamics and relative abundance of three plant species (Antennaria dioica, Hieracum pilosella and Hypochaeris maculata) in dry semi-natural grasslands', *Ecography* **20**(6), 559 - 568.
- (107) Eriksson, O. (1996), 'Population ecology and conservation: Some theoretical considerations with examples from the Nordic flora', *Acta Universitatis Upsaliensis Symbolae Botanicae Upsalienses* **31**(3), 159--167.
- (108) Eriksson, O. & Kiviniemi, K. (1999), 'Site occupancy, recruitment and extinction thresholds in grassland plants: an experimental study', *Biological Conservation* **87**(3), 319 - 325.
- (109) Erschbamer, B.; Mallaun, M. & Unterluggauer, P. (2006), 'Plant diversity along altitudinal gradients in the Southern and Central Alps of South Tyrol and Trentino (Italy)', *Gredleriana* **6**, 47--68.
- (110) Escudero, M.; Valcárcel, V.; Vargas, P. & Luceño, M. (2010), 'Bipolar disjunctions in Carex: Long-distance dispersal, vicariance, or parallel evolution?', *Flora - Morphology, Distribution, Functional Ecology of Plants* **205**(2), 118 - 127.
- (111) Essl, F. (2005), 'Remarkable floristic records from Vienna, Lower Austria, Burgenland and Syria, Part II', *Linzer Biologische Beitrage* **37**(2), 1207--1230.
- (112) Essl, F. (2003), 'Remarkable floristic records from Vienna, Lower Austria, Burgenland and Styria.', *Linzer Biologische Beitrage* **35**(2), 935--956.
- (113) Eysink, F. & De Bruijn, O. (1997), 'Restoration of fen meadows in the Lemselermaten', *Levende Natuur* **98**(7), 258--265.
- (114) Fadeeva, I. A. & Bogomolova, T. V. (2006), 'Records of rare plants in Smolensk province', *Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody Otdel Biologicheskii* **111**(Part 6), 67.
- (115) de Faria, A. L. (1997), 'The atomic basis of biological symmetry and periodicity', *Biosystems* **43**(2), 115 - 135.
- (116) Feldt, C. (2008), 'Beiträge zur Populationsdynamik von Antennaria dioica (L.) Gaertn. im Bereich der mittleren Schwäbischen Alb. Dissertation', PhD thesis, Universität Ulm.
- (117) Ferrari, C.; Bonafede, F. & Alessandrini, A. (1993), 'Rare plants of the Emilia-Romagna region

- (Northern Italy): A data bank and computer-mapped atlas for conservation purposes', *Biological Conservation* **64**(1), 11 - 18.
- (118) FIASSON, J. L.; GLUCHOFFFIASSON, K.; MUGNIER, C.; BARGHI, N. & SILJAKYAKOVLEV, S. (1991), 'Flavonoid Analysis of European Species of the Genus *Hypochoeris* (asteraceae)', *Biochemical Systematics and Ecology* **19**(2), 157--162.
- (119) Fink, S. (1996), 'Verbreitung, Soziologie und Ökologie von *Pedicularis palustris* L. in Nordwestdeutschland', Master's thesis, Universität Bremen.
- (120) Fischer, M. & Wipf, S. (2002), 'Effect of low-intensity grazing on the species-rich vegetation of traditionally mown subalpine meadows', *Biological Conservation* **104**(1), 1 - 11.
- (121) Franz, G. (1979), 'Metabolism of Reserve Polysaccharides in Tubers of *Orchis morio* L', *Planta Med* **36**(5), 68-73.
- (122) Franz, G. (1973), 'Biosynthesis of salep mannan', *Phytochemistry* **12**(10), 2369 - 2373.
- (123) Franzen, D. & Eriksson, O. (2003), 'Patch distribution and dispersal limitation of four plant species in Swedish semi-natural grasslands', *Plant Ecology* **166**(2), 217--225.
- (124) Fyfe, R. M.; Brown, A. G. & Rippon, S. J. (2004), 'Characterising the late prehistoric, [] Romano-British' and medieval landscape, and dating the emergence of a regionally distinct agricultural system in South West Britain', *Journal of Archaeological Science* **31**(12), 1699 - 1714.
- (125) GABRIELIAN, E. T. & GAMBARYAN, P. P. (1973), 'New and Rare Floristic Finds In the Armenian-ssr', *Biologicheskii Zhurnal Armenii* **26**(11), 56--60.
- (126) Garcia-Baquero Moneo, G. & Valle Gutierrez, C. J. (1999), 'New plant species in the Sierra de la Demanda (La Rioja, Spain)', *Studia Botanica* **18**, 145--148.
- (127) Gattiker, H. (1938), 'Vergleichende anatomisch-pharmakognostische Untersuchung einiger Drogen der Gattungen *Antennaria*, *Gnaphalium* und *Helichrysum*', PhD thesis, ETH Zürich.
- (128) Gawalowski, G. & Karrer, G. (2001), 'On the growth form of *Phyteuma ovatum* (Campanulaceae)', *Neilreichia* **1**, 165--175.
- (129) Gigord, L. D. B.; Macnair, M. R. & Smithson, A. (2001), 'Negative frequency-dependent selection maintains a dramatic flower color polymorphism in the rewardless orchid *Dactylorhiza sambucina* (L.) Soo`', *PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA* **98**, 6253-6255.
- (130) Gillman, M. & Dodd, M. (1998), 'The variability of orchid population size', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* **126**(1-2), 65-74.
- (131) Gillman, M. P. & Dodd, M. (2000), 'Detection of delayed density dependence in an orchid population', *Journal of Ecology* **88**(2), 204--212.

- (132) gemeinnützige GmbH, N. A. (), 'Wildwachsende Orchideen im Landkreis Annaberg'.
- (133) Godefroid, S. (2001), 'Temporal analysis of the Brussels flora as indicator for changing environmental quality', *Landscape and Urban Planning* **52**(4), 203 - 224.
- (134) Gorissen, I. (1992), 'Noteworthy occurrences of *Platanthera bifolia*, *Azolla filliculoides*, and other species at the southern edge of the Lower Rhine Basin', *Floristische Rundbriefe* **26**(1), 21-22.
- (135) Gostin, I. N. (2009), 'Air Pollution Effects on the Leaf Structure of some Fabaceae Species', *Notulae Botanicae Horti Agrobotanici Cluj-napoca* **37**(2), 57--63.
- (136) Grabherr, G. (2009), 'Biodiversity in the high ranges of the Alps: Ethnobotanical and climate change perspectives', *Global Environmental Change* **19**(2), 167 - 172.
- (137) Grayer, R. J. & Kokubun, T. (2001), 'Plant-fungal interactions: the search for phytoalexins and other antifungal compounds from higher plants', *Phytochemistry* **56**(3), 253 - 263.
- (138) GROOTJANS, A. P. (1979), 'Some Remarks On the Relation Between Nitrogen Mineralization, Groundwater Table and Standing Crop In Wet Meadows', *Acta Botanica Neerlandica* **28**(2-3), 234--235.
- (139) Grubb, P. J. (1998), 'A reassessment of the strategies of plants which cope with shortages of resources', *Perspectives in Plant Ecology, Evolution and Systematics* **1**(1), 3 - 31.
- (140) Gruber, M. (1984), 'The Marshes With *Eriophorum latifolium* and Sedges In the Valley of Louron Central Pyrenees France', *Bulletin de la Societe d'Histoire Naturelle de Toulouse* **120**, 27-30.
- (141) Grünanger, P.; Caporali, E.; Marziani, G.; Menguzzato, E. & Servettaz, O. (1998), 'Molecular (RAPD) analysis on Italian taxa of the *Ophrys bertolonii* aggregate (Orchidaceae)', *PLANT SYSTEMATICS AND EVOLUTION* **212**(3-4), 177-184.
- (142) Gucwa-Przepiora, E. & Blaszkowski, J. (2007), 'Arbuscular mycorrhiza of plants spontaneously colonizing the soda heap in Jaworzno (Southern Poland)', *Acta Societatis Botanicorum Poloniae* **76**(1), 69--74.
- (143) Gugerli, F. & Holderegger, R. (2001), 'Nunatak survival, tabula rasa and the influence of the Pleistocene ice-ages on plant evolution in mountain areas', *Trends in Plant Science* **6**(9), 397--398.
- (144) Gustavsson, E.; Lennartsson, T. & Emanuelsson, M. (2007), 'Land use more than 200 years ago explains current grassland plant diversity in a Swedish agricultural landscape', *Biological Conservation* **138**(1-2), 47 - 59.
- (145) Gutermann, W. (2009), 'Notulae nomenclaturales 29-40', *Phyton-annales Rei Botanicae* **49**(1), 77--92.

- (146) H. F. Beyrle, S. E. Smith, C. M. M. F. R. L. P. (1995), 'Colonization of *Orchis morio* protocorms by a mycorrhizal fungus: effects of nitrogen nutrition and glyphosate in modifying the responses', *Canadian Journal of Botany* **73**, 1128-1140.
- (147) H. Kretzschmar, W. Eccarius, H. D. (2007), *Die Orchideengattungen Anacamptis, Orchis, Neotinea Phylogenie, Taxonomie, Morphologie, Biologie, Verbreitung, Ökologie und Hybridisation*, Echinomedia.
- (148) Haavik, S.; Paulsen, B. S.; Wold, J. K. & Øivind Grimme (1982), 'Arabinogalactans in a purified allergen preparation from pollen of timothy (*Phleum pratense*)', *Phytochemistry* **21**(8), 1913 - 1919.
- (149) Hackel, E. (1883), 'Über das Vorkommen von *Calamagrostis phragmitioides* (HARTM.) in Deutschland', *Bericht Deutsche Botanische Gesellschaft* **1**, 161-162.
- (150) Hackel, E. (1883), 'Über das Vorkommen von *Calamagrostis phragmitioides* (HARTM.) in Deutschland', *Bericht Deutsche Botanische Gesellschaft* **1**, 161-162.
- (151) Harastova-Sobotkova, M.; Jersakova, J.; Kindlmann, P. & Curn, L. (2005), 'Morphometric and genetic divergence among populations of *Neotinea ustulata* (Orchidaceae) with different flowering phenologies', *FOLIA GEOBOTANICA* **40**(4), 385-405.
- (152) Heckford, R. J. (2010), '*Scrobipalpa murinella* (Duponchel, 1843) (Lepidoptera: Gelechiidae) in Northern Ireland, and a note on the larva', *Entomologist's Gazette* **61**(2), 132.
- (153) Heckford, R. J. (2001), '*Coleophora pappiferella* Hofmann, 1869 (Lepidoptera: Coleophoridae) new to Britain and *Scrobipalpa murinella* (Duponchel, 1843) (Lepidoptera: Gelechiidae) at the same locality', *Entomologist's Gazette* **52**(2), 101--104.
- (154) Heckford, R. J. (2001), 'Observations on the larval habits of *Kessleria saxifragae* (Stainton) (Lepidoptera: Yponomeutidae)', *Entomologist's Gazette* **52**(1), 49--50.
- (155) Hedberg, P. & Kotowski, W. (2010), 'New nature by sowing? The current state of species introduction in grassland restoration, and the road ahead', *Journal for Nature Conservation* **18**(4), 304 - 308.
- (156) Heggland, A. (2005), 'New findings of lanceleaf grapefern *Botrychium lanceolatum* in Tinn, Telemark', *Blyttia* **63**(4), 213--215.
- (157) HEIJNE, B.; HOFSTRA, J. J.; HEIL, G. W.; VANDAM, D. & BOBBINK, R. (1992), 'Effect of the Air-pollution Component Ammonium-sulfate On the Vam Infection-rate of 3 Heathland Species', *Plant and Soil* **144**(1), 1--12.
- (158) Heinrich, W. & Dietrich, H. (2008), 'Heimische Orchideen in urbanen Biotopen', *Feddes Repert.* **119**(5-6), 388--432.
- (159) Hejcman, M.; Schellberg, J. & Pavlu, V. (2010), '*Dactylorhiza maculata*, *Platanthera bifolia* and

- Listera ovata survive N application under P limitation', *Acta Oecologica In Press, Corrected Proof*, - .
- (160) Hempel, W. (1982), 'Über das Indigenat von *Carlina acaulis* am Pöhlberg bei Annaberg', *Gled* **9**, 201-203.
- (161) Herzog, F.; Dreier, S.; Hofer, G.; Marfurt, C.; Schüpbach, B.; Spiess, M. & Walter, T. (2005), 'Effect of ecological compensation areas on floristic and breeding bird diversity in Swiss agricultural landscapes', *Agriculture, Ecosystems & Environment* **108**(3), 189 - 204.
- (162) Hiemeyer, F. (1991), 'Orobanche-elatior Sutton In Ries', *Berichte der Bayerischen Botanischen Gesellschaft zur Erforschung der Heimischen Flora* **62**, 270--271.
- (163) Hodgson, J.; Grime, J.; Wilson, P.; Thompson, K. & Band, S. (2005), 'The impacts of agricultural change (1963-2003) on the grassland flora of Central England: processes and prospects', *Basic and Applied Ecology* **6**(2), 107 - 118.
- (164) Hogstrom, S. (1995), 'Prunella laciniata in Gotland, Sweden', *Svensk Botanisk Tidskrift* **89**(6), 333--334.
- (165) Honnay, O.; Hermy, M. & Coppin, P. (1999), 'Impact of habitat quality on forest plant species colonization', *Forest Ecology and Management* **115**(2-3), 157 - 170.
- (166) Honnay, O.; Hermy, M. & Coppin, P. (1999), 'Effects of area, age and diversity of forest patches in Belgium on plant species richness, and implications for conservation and reforestation', *Biological Conservation* **87**(1), 73 - 84.
- (167) Hopper, S. D. & Lambers, H. (2009), 'Darwin as a plant scientist: a Southern Hemisphere perspective', *Trends in Plant Science* **14**(8), 421 - 435.
- (168) Hossain, M. M. (2010), 'Therapeutic Orchids: Traditional Uses and Recent Advances - An Overview', *Fitoterapia In Press, Accepted Manuscript*, - .
- (169) Hrevusová, Z.; Hejzman, M.; Pavlu, V. V.; Hakl, J.; Klaudivová, M. & Mrkvicka, J. (2009), 'Long-term dynamics of biomass production, soil chemical properties and plant species composition of alluvial grassland after the cessation of fertilizer application in the Czech Republic', *Agriculture, Ecosystems & Environment* **130**(3-4), 123 - 130.
- (170) Hrivnak, R. & Ujhazy, K. (2005), 'Changes of the mountain grassland vegetation after abandonment and colonization by Norway spruce', *Ekologia-bratislava* **24**(3), 231--253.
- (171) Huelber, K.; Sonnleitner, M.; Flatscher, R.; Berger, A.; Dobrovsky, R.; Niessner, S.; Nigl, T.; Schneeweiss, G. M.; Kubesova, M.; Rauchova, J.; Suda, J. & Schoenswetter, P. (2009), 'Ecological segregation drives fine-scale cytotype distribution of *Senecio carniolicus* in the Eastern Alps', *Preslia (Prague)* **81**(3), 309--319.
- (172) Huhta, A. P.; Hellstrom, K.; Rautio, P. & Tuomi, J. (2003), 'Grazing tolerance of *Gentianella*

- amarella and other monocarpic herbs: why is tolerance highest at low damage levels?', *Plant Ecology* **166**(1), 49-61.
- (173) Hurst, A. & John, E. (1999), 'The biotic and abiotic changes associated with *Brachypodium pinnatum* dominance in chalk grassland in south-east England', *Biological Conservation* **88**(1), 75 - 84.
- (174) Hylgaard, T. (1980), 'Recovery of plant communities on coastal sand-dunes disturbed by human trampling', *Biological Conservation* **19**(1), 15 - 25.
- (175) Hájek, M.; Horsák, M.; Hájková, P. & Díte, D. (2006), 'Habitat diversity of central European fens in relation to environmental gradients and an effort to standardise fen terminology in ecological studies', *Perspectives in Plant Ecology, Evolution and Systematics* **8**(2), 97 - 114.
- (176) Ihse, M. & Lindahl, C. (2000), 'A holistic model for landscape ecology in practice: the Swedish survey and management of ancient meadows and pastures', *Landscape and Urban Planning* **50**(1-3), 59 - 84.
- (177) Inghe, O. & Tamm, C. O. (1988), 'Survival and Flowering of Perennial Herbs. V Patterns of Flowering', *Oikos* **51**, 203-219.
- (178) Inghe, O. & Tamm, C. O. (1988), 'Survival and flowering of perennial herbs. V. Patterns of flowering', *Oikos* **51**(2), 203-219.
- (179) Irving, A. (1875), 'The Geology of the Nottingham District.–Part II', *Proceedings of the Geologists' Association* **4**(1), 57 - 90.
- (180) Jacquemyn, H.; Brys, R.; Adriaens, D.; Honnay, O. & Roldan-Ruiz, I. (2009), 'Effects of population size and forest management on genetic diversity and structure of the tuberous orchid *Orchis mascula*', *CONSERVATION GENETICS* **10**(1), 161-168.
- (181) Jacquemyn, H.; Brys, R.; Hermy, M. & Willems, J. H. (2005), 'Does nectar reward affect rarity and extinction probabilities of orchid species? An assessment using historical records from Belgium and the Netherlands', *Biological Conservation* **121**(2), 257 - 263.
- (182) Jacquemyn, H.; Brys, R.; Honnay, O. & Hermy, M. (2008), 'Effects of coppicing on demographic structure, fruit and seed set in *Orchis mascula*', *BASIC AND APPLIED ECOLOGY* **9**(4), 392-400.
- (183) Jacquemyn, H.; Brys, R.; Honnay, O. & Hutchings, M. J. (2009), 'Biological Flora of the British Isles: *Orchis mascula* (L.) L.', *Journal of Ecology* **97**(2), 360--377.
- (184) Jacquemyn, H.; Honnay, O.; Cammue, B. P. A.; Brys, R. & Lievens, B. (2010), 'Low specificity and nested subset structure characterize mycorrhizal associations in five closely related species of the genus *Orchis*', *MOLECULAR ECOLOGY* **19**(18), 4086-4095.
- (185) Jacquemyn, H.; Vandepitte, K.; Brys, R.; Honnay, O. & Roldan-Ruiz, I. (2007), 'Fitness variation and genetic diversity in small, remnant populations of the food deceptive orchid *Orchis*

- purpurea', *BIOLOGICAL CONSERVATION* **139**(1-2), 203-210.
- (186) Jacquemyn, H.; Wiegand, T.; Vandepitte, K.; Brys, R.; Roldán-Ruiz, I. & Honnay, O. (2009), 'Multigenerational analysis of spatial structure in the terrestrial, food-deceptive orchid *Orchis mascula*', *Journal of Ecology* **97**(2), 206--216.
- (187) Janecek, S. & Leps, J. (2005), 'Effect of litter, leaf cover and cover of basal internodes of the dominant species *Molinia caerulea* on seedling recruitment and established vegetation', *Acta Oecologica* **28**(2), 141 - 147.
- (188) Janecková, P.; Wotavová, K.; Schödelbauerová, I.; Jersáková, J. & Kindlmann, P. (2006), 'Relative effects of management and environmental conditions on performance and survival of populations of a terrestrial orchid, *Dactylorhiza majalis*', *Biological Conservation* **129**(1), 40 - 49.
- (189) Jankovská, V. & Rybníček, K. (1988), 'The genus *Carex* in the late glacial and holocene of Czechoslovakia', *Aquatic Botany* **30**(1-2), 23 - 37.
- (190) Jansen, A. J. M.; Grootjans, A. P. & Jalink, M. H. (2000), 'Hydrology of Dutch *Cirsio-Molinietum* meadows: Prospects for restoration', *Applied Vegetation Science* **3**(1), 51--64.
- (191) Jantunen, J. & Saarinen, K. (2002), 'Vegetation of uncultivated arable land under different management in Finnish and Russian Karelia', *Agriculture, Ecosystems & Environment* **92**(2-3), 225 - 233.
- (192) Jersakova, J. & Johnson, S. (2006), 'Lack of floral nectar reduces self-pollination in a fly-pollinated orchid', *OECOLOGIA* **147**(1), 60-68.
- (193) Jersakova, J. & Kindlmann, P. (1998), 'Patterns of pollinator-generated fruit set in *Orchis morio* (Orchidaceae)', *FOLIA GEOBOTANICA* **33**(4), 377-390.
- (194) Jersakova, J.; Kindlmann, P. & Renner, S. S. (2006), 'Is the colour dimorphism in *Dactylorhiza sambucina* maintained by differential seed viability instead of frequency-dependent selection?', *Folia Geobotanica* **41**(1), 61-76.
- (195) Jersáková, J.; Johnson, S. D.; Kindlmann, P. & Pupin, A.-C. (2008), 'Effect of nectar supplementation on male and female components of pollination success in the deceptive orchid *Dactylorhiza sambucina*', *Acta Oecologica* **33**(3), 300 - 306.
- (196) Johansson, K.-A. (2009), 'Journey to Romania', *Svensk Botanisk Tidskrift* **103**(6), 323--329.
- (197) Johnson, M.; Putwain, P. & Holliday, R. (1978), 'Wildlife conservation value of derelict metalliferous mine workings in Wales', *Biological Conservation* **14**(2), 131 - 148.
- (198) Johnson, S. & Nilsson, L. (1999), 'Pollen carryover, geitonogamy, and the evolution of deceptive pollination systems in orchids', *ECOLOGY* **80**(8), 2607-2619.

- (199) Johnson, S.; Peter, C. & Agren, J. (2004), 'The effects of nectar addition on pollen removal and geitonogamy in the non-rewarding orchid *Anacamptis morio*', *PROCEEDINGS OF THE ROYAL SOCIETY OF LONDON SERIES B-BIOLOGICAL SCIENCES* **271**(1541), 803-809.
- (200) Johnson, S.; Peter, C.; Nilsson, L. & Agren, J. (2003), 'Pollination success in a deceptive orchid is enhanced by co-occurring rewarding magnet plants', *ECOLOGY* **84**(11), 2919-2927.
- (201) JONASSON, S. & CALLAGHAN, T. V. (1992), 'Root Mechanical-properties Related To Disturbed and Stressed Habitats In the Arctic', *New Phytologist* **122**(1), 179--186.
- (202) Jones, E. & Hughes, R. E. (1983), 'Foliar ascorbic acid in some angiosperms', *Phytochemistry* **22**(11), 2493 - 2499.
- (203) Jongepierová, I.; Mitchley, J. & Tzanopoulos, J. (2007), 'A field experiment to recreate species rich hay meadows using regional seed mixtures', *Biological Conservation* **139**(3-4), 297 - 305.
- (204) Juillet, N.; Dunand-Martin, S. & Gigord, L. D. B. (2007), 'Evidence for inbreeding depression in the food-deceptive colour-dimorphic orchid *Dactylorhiza sambucina* (L.) Soo', *PLANT BIOLOGY* **9**(1), 147-151.
- (205) Juillet, N.; Dunand-Martin, S. & Gigord, L. D. B. (2007), 'Evidence for inbreeding depression in the food-deceptive colour-dimorphic orchid *Dactylorhiza sambucina* (L.) Soo', *Plant Biology* **9**(1), 147-151.
- (206) Juillet, N. & Scopece, G. (2010), 'Does floral trait variability enhance reproductive success in deceptive orchids?', *Perspectives in Plant Ecology, Evolution and Systematics* **12**(4), 317 - 322.
- (207) Jurzyk, S. & Wrobel, M. (2003), 'Co-occurrence of two species *Molinia caerulea* L. and "red-list" species *Carex pulicaris* L. in Western Pomerania (Poland)', *Polish Journal of Ecology* **51**(3), 363--367.
- (208) KAMYSHEV, N. S. (1979), 'Comparative Characteristics of Dry Meadows and Caespitose Herb Steppes', *Biologicheskie Nauki (Moscow)*(3), 62--66.
- (209) Karrenberg, S. & Jensen, K. (2000), 'Effects of pollination and pollen source on the seed set of *Pedicularis palustris*', *Folia Geobotanica* **35**(2), 191-202.
- (210) KAZAKOV, A. Z. (1977), 'Flavonoids of *Trifolium-montanum*', *Khimiya Prirodnikh Soedinenii*(3), 415 - 416.
- (211) KAZIMIERSKA, E. M. & KAZIMIERSKI, T. (1985), 'Crossing of Species In the Genus *Trifolium* L .1. *Trifolium-repens* Lxt-*montanum* L - Fertilization and Development of the Embryo and Endosperm', *Genetica Polonica* **26**(3), 327--.
- (212) Kelly, D. (1989), 'Demography of Short-lived Plants In Chalk Grassland: 1. Life-cycle Variation In Annuals and Strict Biennials', *Journal of Ecology* **77**(3), 747-769.

- (213) Kelly, D. (1989), 'Demography of Short-lived Plants In Chalk Grassland: 2. Control of Mortality and Fecundity', *Journal of Ecology* **77**(3), 770-784.
- (214) Kelly, D. (1989), 'Demography of Short-lived Plants In Chalk Grassland: 3. Population Stability', *Journal of Ecology* **77**(3), 785-798.
- (215) Kepczynski, K. & Cyzman, W. (1992), 'Notable flora of the forest district Grodno', *Acta Universitatis Nicolai Copernici Biologia* **40**(0), 3--58.
- (216) Kindlmann, P. & Jersáková, J. (2006-03-01), 'Effect of floral display on reproductive success in terrestrial orchids', *Folia Geobotanica* **41**(1), 47--60.
- (217) Kirichok, E. I. (1999), 'The record of *Swertia perennis* (Gentianaceae) northwest of Smolensk region', *Botanicheskii Zhurnal (St. Petersburg)* **84**(2), 124-127.
- (218) Kirschner, J.; Kirschnerova, L. & Stepanek, J. (2007), 'Generally accepted plant names based on material from the Czech Republic and published in 1753-1820', *Preslia* **79**(4), 323--365.
- (219) Klanderud, K. (2010), 'Species recruitment in alpine plant communities: the role of species interactions and productivity', *Journal of Ecology* **98**(5), 1128--1133.
- (220) Kleijn, D.; Bekker, R. M.; Bobbink, R.; De Graaf, M. C. C. & Roelofs, J. G. M. (2008), 'In search for key biogeochemical factors affecting plant species persistence in heathland and acidic grasslands: a comparison of common and rare species', *Journal of Applied Ecology* **45**(2), 680--687.
- (221) Klein, E. (2000), 'Pseudorchis albida subsp. *tricuspis* (BECK) KLEIN stat. nov., a largely overlooked calcicolous, alpic-boreal taxon (Orchidaceae-Orchideae)', *PHYTON-ANNALES REI BOTANICAE* **40**(1), 141-159
- (222) Klett, J. E.; Staats, D. & Rogoyski, M. (2005), 'Preemergence weed control in container-grown herbaceous perennials', *Hortscience* **40**(4), 1024.
- (223) Klimes[breve]ová, J. & Klimes, L. (2007), 'Bud banks and their role in vegetative regeneration - A literature review and proposal for simple classification and assessment', *Perspectives in Plant Ecology, Evolution and Systematics* **8**(3), 115 - 129.
- (224) Klinkhamer, P. G. L.; De Jong, T. J. & Meelis, E. (1991), 'The control of flowering in the monocarpic perennial *Carlina vulgaris*', *Oikos* **61**(1), 88-95.
- (225) Knudsen, J. T.; Tollsten, L. & Bergström, L. (1993), 'Floral scents—a checklist of volatile compounds isolated by head-space techniques', *Phytochemistry* **33**(2), 253 - 280.
- (226) Koch, M. A.; Scheriau, C.; Schupfner, M. & Bernhardt, K.-G. (2010), 'Long-term monitoring of the restoration and development of limestone grasslands in north western Germany: Vegetation screening and soil seed bank analysis', *Flora - Morphology, Distribution, Functional Ecology of Plants* **In Press, Corrected Proof**, - .

- (227) Koeman-Kwak, M. (1973), 'Pollination of Pedicularis-palustris By Nectar Thieves (short-tongued Bumblebees)', *Acta Botanica Neerlandica* **22**(6), 608-615.
- (228) Kohn, D. (2005), 'The miraculous season', *NATURAL HISTORY* **114**(9), 38-40.
- (229) Kolb, A. (2008), 'Habitat fragmentation reduces plant fitness by disturbing pollination and modifying response to herbivory', *Biological Conservation* **141**(10), 2540--2549.
- (230) Kolb, A. (2005), 'Reduced reproductive success and offspring survival in fragmented populations of the forest herb *Phyteuma spicatum*', *Journal of Ecology* **93**(6), 1226--1237.
- (231) Kolb, A. & Barsch, K. (2010), 'Environmental factors and seed abundance influence seedling emergence of a perennial forest herb', *Acta Oecologica* **36**(5).
- (232) Kolb, A.; Dahlgren, J. P. & Ehrlen, J. (2010), 'Population size affects vital rates but not population growth rate of a perennial plant', *Ecology* **91**(11), 3210--3217.
- (233) Krause, A. & Wolf, G. (1991), 'Das Purpureitgras (*Calamagrostis purpurea* (Trin.) Trin. ssp. *phragmitoides* (Hartm.) Tzvelev) neu für Nordrhein-Westfalen', *Dechenia* **144**, 115.
- (234) Kristiansen, K. A.; Freudenstein, J. V.; Rasmussen, F. N. & Rasmussen, H. N. (2004), 'Molecular identification of mycorrhizal fungi in *Neuwiedia veratrifolia* (Orchidaceae)', *Molecular Phylogenetics and Evolution* **33**(2), 251 - 258.
- (235) KROEZE, C.; PEGTEL, D. M. & BLOM, C. J. C. (1989), 'An Experimental Comparison of Aluminum and Manganese Susceptibility In *Antennaria-dioica*, *Arnica-montana*, *Viola-canina*, *Filago-minima* and *Deschampsia-flexuosa*', *Acta Botanica Neerlandica* **38**(2), 165--172.
- (236) Kropf, M. (1995), 'Vegetationskundlicher Vergleich von Standorten des Holunder-Knabenkrautes (*Dactylorhiza sambucina*) im Nahegebiet und der Rheinhessischen Schweiz unter Berücksichtigung von Pflege sowie Verbreitung und Gefährdung dieser Orchideenart.', *Flora Fauna Rheinland-Pfalz* **8**, 133-145.
- (237) Kropf, M. & Renner, S. S. (2008), 'Pollinator-mediated selfing in two deceptive orchids and a review of pollinium tracking studies addressing geitonogamy', *OECOLOGIA* **155**(3), 497-508.
- (238) Kropf, M. & Renner, S. S. (2005), 'Pollination success in monochromic yellow populations of the rewardless orchid *Dactylorhiza sambucina*', *Plant Systematics and Evolution* **254**(3-4), 185-197.
- (239) Kruse, J. (1987), 'Morphological Investigations of Involucral and Receptacular Bracts of the Genus *Carlina* L', *Flora* **179**(1), 17-33.
- (240) Kull, T. & Hutchings, M. J. (2006), 'A comparative analysis of decline in the distribution ranges of orchid species in Estonia and the United Kingdom', *Biological Conservation* **129**(1), 31 - 39.
- (241) Kull, T.; Kindlmann, P.; Hutchings, M. J. & Primack, R. B. (2006), 'Conservation biology of orchids: Introduction to the special issue', *Biological Conservation* **129**(1), 1 - 3.

- (242) Kwak, M. M. (2002), 'Bumblebees as flower visitors: Pollinators and profiteers.', *Entomologische Berichten (Amsterdam)* **62**(3-4), 73--81.
- (243) Kytöviita, M.-M.; Vestberg, M. & Tuomi, J. (2003), 'A test of mutual aid in common mycorrhizal networks: established vegetation negates benefit in seedlings', *Ecology* **84**(4), 898-906.
- (244) Landi, M.; Frignani, F.; Lazzeri, C. & Angiolini, C. (2009), 'Abundance of orchids on calcareous grasslands in relation to community species, environmental, and vegetational conditions', *RUSSIAN JOURNAL OF ECOLOGY* **40**(7), 486-494.
- (245) de Langhe, J.-E. & Delvosalle, L. (1973), 'Calamagrostis-phragmitoides An Indigenous Grass of Belgium', *Bulletin de la Societe Royale de Botanique de Belgique* **106**(1), 61-64.
- (246) Larsson, E. & Örstadius, L. (2008), 'Fourteen coprophilous species of Psathyrella identified in the Nordic countries using morphology and nuclear rDNA sequence data', *Mycological Research* **112**(10), 1165 - 1185.
- (247) Lassueur, T.; Joost, S. & Randin, C. F. (2006), 'Very high resolution digital elevation models: Do they improve models of plant species distribution?', *Ecological Modelling* **198**(1-2), 139 - 153.
- (248) Lauber, W. & Körner, C. (1997), 'In situ stomatal responses to long-term CO₂ enrichment in calcareous grassland plants', *Acta Oecologica* **18**(3), 221 - 229.
- (249) Leck, M. A. & Schütz, W. (2005), 'Regeneration of Cyperaceae, with particular reference to seed ecology and seed banks', *Perspectives in Plant Ecology, Evolution and Systematics* **7**(2), 95 - 133.
- (250) Leins, P. & Erbar, C. (2006), 'Secondary pollen presentation syndromes of the Asterales - a phylogenetic perspective', *Botanische Jahrbuecher fuer Systematik Pflanzengeschichte und Pflanzengeographie* **127**(1), 83--103.
- (251) Leins, P. & Erbar, C. (2003), 'The pollen box in Cyphiaceae (Campanulales)', *International Journal of Plant Sciences* **164**(5), S321--S328.
- (252) Leps, J. (2004), 'What do the biodiversity experiments tell us about consequences of plant species loss in the real world?', *Basic and Applied Ecology* **5**(6), 529 - 534.
- (253) LfULGAbt. Natur, Landschaft, B. L., ed. (2009), *Naturschutzfachliche Aspekte des Grünlandes in Sachsen*, Landesamt für Umwelt, Landwirtschaft und Geologie.
- (254) Lienert, J.; Diemer, M. & Schmid, B. (2002), 'Effects of habitat fragmentation on population structure and fitness components of the wetland specialist *Swertia perennis* L. (Gentianaceae)', *Basic and Applied Ecology* **3**(2), 101-114.
- (255) Lienert, J. & Fischer, M. (2004), 'Experimental inbreeding reduces seed production and germination independent of fragmentation of populations of *Swertia perennis*', *Basic and Applied Ecology* **5**(1), 43-52.

- (256) Lienert, J.; Fischer, M. & Diemer, M. (2002), 'Local extinctions of the wetland specialist *Swertia perennis* L. (Gentianaceae) in Switzerland: A revisitation study based on herbarium records', *Biological Conservation* **103**(1), 65-76.
- (257) Lienert, J.; Fischer, M.; Schneller, J. & Diemer, M. (2002), 'Isozyme variability of the wetland specialist *Swertia perennis* (Gentianaceae) in relation to habitat size, isolation, and plant fitness', *American Journal of Botany* **89**(5), 801-811.
- (258) Lievens, B.; van Kerckhove, S.; Justé, A.; Cammue, B. P.; Honnay, O. & Jacquemyn, H. (2010), 'From extensive clone libraries to comprehensive DNA arrays for the efficient and simultaneous detection and identification of orchid mycorrhizal fungi', *Journal of Microbiological Methods* **80**(1), 76 - 85.
- (259) van der Linden, M.; Vickery, E.; Charman, D. J.; Borekens, P. & van Geel, B. (2008), 'Vegetation history and human impact during the last 300 years recorded in a German peat deposit', *Review of Palaeobotany and Palynology* **152**(3-4), 158--175.
- (260) van der Linden, M.; Vickery, E.; Charman, D. J.; Broekens, P. & van Geel, B. (2008), 'Vegetation history and human impact during the last 300 years recorded in a German peat deposit', *Review of Palaeobotany and Palynology* **152**(3-4), 158 - 175.
- (261) Lippert, W. (1992), 'Contributions to the floristic mapping of Bavaria: A supplement to the Bavarian Atlas', *Berichte der Bayerischen Botanischen Gesellschaft zur Erforschung der Heimischen Flora* **0**(SUPPL. 5), 3--64.
- (262) Llamas, F.; Acedo, C.; Andres, J.; Carbo, R.; Perez, M. A. & Penas, A. (1993), 'Anatomy of the European species of *Carex* L. Sect. *Unciniiformes* Kuk. (Cyperaceae)', *Anales del Jardin Botanico de Madrid* **51**(1), 97--109.
- (263) Lombini, A.; Dinelli, E.; Ferrari, C. & Simoni, A. (1998), 'Plant-soil relationships in the serpentinite screes of Mt. Prinzera (Northern Apennines, Italy)', *Journal of Geochemical Exploration* **64**(1-3), 19 - 33.
- (264) Ludwig, W. (1991), 'Purple Reed Grass *Calamagrostis-phragmitoides* In Hesse and Adjacent Regions', *Hessische Floristische Briefe* **40**(2), 17-20.
- (265) Lutz, M.; Piatek, M.; Kemler, M.; Chlebicki, A. & Oberwinkler, F. (2008), 'Anther smuts of Caryophyllaceae: Molecular analyses reveal further new species', *Mycological Research* **112**(11), 1280 - 1296.
- (266) LYUBARSKII, E. L.; POLUYANOVA, V. I. & SILINA, L. G. (1975), 'Seasonal Variations of the Morphological Structure of Community Populations of Mouse-ear Hawkweed and Common Cats-ear', *Biologicheskije Nauki (Moscow)* **18**(10), 75--79.
- (267) M. Allimuthu, T. M. (2009), 'The Role of *Salamisri* (*Orchis mascula*) in Geriatric Care.', *Siddha Papers* **1**, 1-9.

- (268) Maccherini, S.; Bacaro, G.; Favilli, L.; Piazzini, S.; Santi, E. & Marignani, M. (2009), 'Congruence among vascular plants and butterflies in the evaluation of grassland restoration success', *Acta Oecologica* **35**(2), 311 - 317.
- (269) Macior, L. W. (1993), 'Pollination ecology of *Pedicularis palustris* L. (Scrophulariaceae) in North America', *Plant Species Biology* **8**(1), 35-44.
- (270) Majetic, C.; Rausher, M. & Raguso, R. (2010), 'The pigment-scent connection: Do mutations in regulatory vs. structural anthocyanin genes differentially alter floral scent production in *Ipomoea purpurea*?', *South African Journal of Botany* **76**(4), 632 - 642.
- (271) Ülo Mander; Mikk, M. & Külvik, M. (1999), 'Ecological and low intensity agriculture as contributors to landscape and biological diversity', *Landscape and Urban Planning* **46**(1-3), 169 - 177.
- (272) Marini, L.; Scotton, M.; Klimek, S.; Isselstein, J. & Pecile, A. (2007), 'Effects of local factors on plant species richness and composition of Alpine meadows', *Agriculture, Ecosystems & Environment* **119**(3-4), 281 - 288.
- (273) Marion, B.; Bonis, A. & Bouzillat, J.-B. (2010), 'How Much does Grazing-Induced Heterogeneity Impact Plant Diversity in Wet Grasslands?', *Ecoscience* **17**(3), 229--239.
- (274) Maskova, Z.; Zemek, F. & Kvet, J. (2008), 'Normalized difference vegetation index (NDVI) in the management of mountain meadows', *Boreal Environment Research* **13**(5), 417--432.
- (275) Masuhara, G.; Katsuya, K. & Yamaguchi, K. (1993), 'Potential for symbiosis of *Rhizoctonia solani* and binucleate *Rhizoctonia* with seeds of *Spiranthes sinensis* var. *amoena* in vitro', *Mycological Research* **97**(6), 746 - 752.
- (276) Matthies, D.; Brauer, I.; Maibom, W. & Tschardt, T. (2004), 'Population size and the risk of local extinction: empirical evidence from rare plants', *Oikos* **105**(3), 481--488.
- (277) Mattiason, G. (1986), 'The Decline of *Dactylorhiza-sambucina* In the Nature Reserve Kungsmarken Southernmost Sweden', *Svensk Botanisk Tidskrift* **80**(1), 17-25.
- (278) Maurer, K.; Durka, W. & Stöcklin, J. (2003), 'Frequency of plant species in remnants of calcareous grassland and their dispersal and persistence characteristics', *Basic and Applied Ecology* **4**(4), 307 - 316.
- (279) McKendrick, S. (1996), 'The effects of shade on seedlings of *Orchis morio* and *Dactylorhiza fuchsii* in chalk and clay soil', *NEW PHYTOLOGIST* **134**(2), 343-352.
- (280) McKendrick, S. L. (1996), 'The effects of fertilizer and root competition on seedlings of *Orchis morio* and *Dactylorhiza fuchsii* in chalk and clay soil', *New Phytologist* **134**(2), 335-342.
- (281) Meinunger, L. (1977), 'Purpureitgras, *Calamagrostis phragmitoides* Hartman, in Thuringen und Sachsen gefunden.', *Hercynia* **14**, 53-54.

- (282) Menkovic, N.; Savikin, K.; Tasic, S.; Zdunic, G.; Stesevic, D.; Milosavljevic, S. & Vincek, D. (2010), 'Ethnobotanical study on traditional uses of wild medicinal plants in Prokletije Mountains (Montenegro)', *Journal of Ethnopharmacology In Press, Corrected Proof*, - .
- (283) Menzano, A.; Rambozzi, L. & Rossi, L. (2007), 'A severe episode of wildlife-derived scabies in domestic goats in Italy', *Small Ruminant Research* **70**(2-3), 154 - 158.
- (284) Mericli, A. H. (1983), 'Constituents of *Antennaria dioica*', *Journal of Natural Products* **46**(6), 941-941.
- (285) van der Merwe, M.; Ericson, L.; Walker, J.; Thrall, P. H. & Burdon, J. J. (2007), 'Evolutionary relationships among species of *Puccinia* and *Uromyces* (Pucciniaceae, Uredinales) inferred from partial protein coding gene phylogenies', *Mycological Research* **111**(2), 163 - 175.
- (286) van der Merwe, M. M.; Walker, J.; Ericson, L. & Burdon, J. J. (2008), 'Coevolution with higher taxonomic host groups within the *Puccinia/Uromyces* rust lineage obscured by host jumps', *Mycological Research* **112**(12), 1387 - 1408.
- (287) MINARIK, J. (1950), '*Antennaria dioica* Gartn., *Gnaphalium dioicum* L.', *Cas Cesk Lek* **63**(9-12), 254--60.
- (288) Moccia, M. D.; Widmer, A. & Cozzolino, S. (2007), 'The strength of reproductive isolation in two hybridizing food-deceptive orchid species', *MOLECULAR ECOLOGY* **16**(14), 2855-2866.
- (289) Montieri, S.; Gaudio, L. & Aceto, S. (2004), 'Isolation of the LFY/FLO homologue in *Orchis italica* and evolutionary analysis in some European orchids', *Gene* **333**, 101 - 109.
- (290) Morier, I.; Guenat, C.; Siegwolf, R.; Vedy, J. C. & Schleppei, P. (2008), 'Dynamics of Atmospheric Nitrogen Deposition in a Temperate Calcareous Forest Soil', *Journal of Environmental Quality* **37**(6), 2012--2021.
- (291) Mortensen, L. M. (1993), 'Effects of ozone on growth of several subalpine plant species', *Norwegian Journal of Agricultural Sciences* **7**(2), 129--138.
- (292) Mueller, J. & Kuhbier, H. (2006), 'History, development and vegetation dynamics of a small calcareous fen in the conservation area Reithbruch near Bremen (NW-Germany).', *Abhandlungen Naturwissenschaftlichen Verein zu Bremen* **46**(1), 101--124.
- (293) MUELLER, W. & SCHUMACHER, W. (1986), 'The Distribution of Rare Species of the Calcareous Flat Moor *Caricion-davallianae* In Eifel West Germany', *Decheniana* **139**, 200--201.
- (294) MUGNIER, C. & SILJAKYAKOVLEV, S. (1987), 'Karyological Study In Some Yugoslavian Populations of *Hypochoeris* (compositae)', *Caryologia* **40**(4), 319--325.
- (295) Muncaciu, S.; Gafta, D.; Cristea, V.; Rosca-Casian, O. & Goia, I. (2010), 'Eco-coenotic conditions and structure of *Trollius europaeus* L. populations in an extrazonal habitat complex (Transylvanian Carpathian foothills)', *Flora - Morphology, Distribution, Functional Ecology of*

Plants **205**(11), 711 - 720.

- (296) Muncaciu, S.; Gafta, D.; Cristea, V.; Rosca-Casian, O. & Goia, I. (2010), 'Eco-coenotic conditions and structure of *Trollius europaeus* L. populations in an extrazonal habitat complex (Transylvanian Carpathian foothills)', *Flora - Morphology, Distribution, Functional Ecology of Plants In Press, Corrected Proof*, - .
- (297) Murton, J. B.; Baker, A.; Bowen, D. Q.; Caseldine, C. J.; Coope, G. R.; Currant, A. P.; Evans, J. G.; Field, M. H.; Green, C. P.; Hatton, J.; Ito, M.; Jones, R. L.; Keen, D. H.; Kerney, M. P.; McEwan, R.; McGregor, D. F. M.; Parish, D.; Robinson, J. E.; Schreve, D. C. & Smart, P. L. (2001), 'A late Middle Pleistocene temperate-periglacial-temperate sequence (Oxygen Isotope Stages 7-5e) near Marsworth, Buckinghamshire, UK', *Quaternary Science Reviews* **20**(18), 1787 - 1825.
- (298) Mückschel, C. & Otte, A. (2003), 'Morphometric parameters: an approach for the indication of environmental conditions on calcareous grassland', *Agriculture, Ecosystems & Environment* **98**(1-3), 213 - 225.
- (299) Müller, F. (2001), 'Verbreitung, Soziologie und Gefährdung von *Carex pulicaris* L. in Sachsen', *Berichte der Arbeitsgemeinschaft Sächsische Botaniker* **NF 18**, 35-48.
- (300) Müller, F. (2001), 'Verbreitung, Soziologie und Gefährdung von *Carex pulicaris* in Sachsen', *Ber. AGsB NF 18*, 35-48.
- (301) Müller-Schneider, P. (1986), *Verbreitungsbiologie der Blütenpflanzen Graubündens*, Vol. 85, Geobotanisches Institut, Zürich.
- (302) NABU (2010), 'GRÜNES BAND SACHSEN Vogtlandkreis', pdf, www.naturerbe.de.
- (303) Nantel, P.; Bouchard, A.; Brouillet, L. & Hay, S. (1998), 'Selection of areas for protecting rare plants with integration of land use conflicts: A case study for the west coast of Newfoundland, Canada', *Biological Conservation* **84**(3), 223 - 234.
- (304) NEILAND, M. & WILCOCK, C. (1995), 'MAXIMIZATION OF REPRODUCTIVE SUCCESS BY EUROPEAN ORCHIDACEAE UNDER CONDITIONS OF INFREQUENT POLLINATION', *PROTOPLASMA* **187**(1-4), 39-48.
- (305) Netien, G. (1994), 'Addendum to the flora of Lyonnais: *Carex buxbaumii* Wahl. and *Carex hartmanii* Cajander', *Bulletin Mensuel de la Societe Linneenne de Lyon* **63**(7), 267--268.
- (306) Neumann, S. M. (1999), 'Populationsbiologische und autökologische Untersuchungen an *Carlina acaulis* ssp. *caulescens* (LAM.) Schlüßer et Martens 1834 im geographischen Vergleich', Master's thesis, Universität Göttingen.
- (307) Nickstadt, A. (2000), 'Populationsbiologie und ökologische Morphologie der Halbtrockenrasenart *Carlina acaulis* L.', Master's thesis, Universität Halle-Wittenberg.
- (308) Nickstadt, A. & Jäger, E. J. (2000), 'Beiträge zur Populationsbiologie der Silberdistel (*Carlina*

- acaulis L.)', *Hercynia* **33**, 245-256.
- (309) NIEMI, A. (1970), 'Occurrences of *Carex-pulicaris*-m On the Hango Peninsula South Finland', *Memoranda Societatis pro Fauna et Flora Fennica* **46**, 39--42.
- (310) Nieschalk, A. & Nieschalk, C. (1976), 'Nachtrag und Berichtigung zu einem Beitrag über *Calamagrostis phragmitoides* Hartman in Nordhessen', *Hessische Floristische Briefe* **25**, 68-69.
- (311) NIKOLAEVA, V. G. (1977), 'Plants Used By Ethnic Groups of the Ussr For Treating Diseases of the Liver and Bile Secretory System', *Rastitel'nye Resursy* **13**(2), 396--403.
- (312) NILSSON, L. (1992), 'ANIMAL POLLINATORS ADJUST PLANT GENDER IN RELATION TO FLORAL DISPLAY - EVIDENCE FROM ORCHIS-MORIO (ORCHIDACEAE)', *EVOLUTIONARY TRENDS IN PLANTS* **6**(1), 33-40.
- (313) Nilsson, L. A. (1983), 'Anthecology of *Orchis mascula* (Orchidaceae)', *Nordic Journal of Botany* **3**(2), 157--179.
- (314) Nilsson, L. A. (1980), 'The Pollination Ecology of *Dactylorhiza-sambucina* (orchidaceae)', *Botaniska Notiser* **133**(3), 367-385.
- (315) Nordal, I. & Wischmann, F. (1987), '*Dactylorhiza sambucina* in Norway [norwegisch]', *Blyttia* **45**(1), 30-38.
- (316) NORDERHAUG, A. (1995), 'Mating Systems of 3 Meadow Plant-species', *Nordic Journal of Botany* **15**(3), 243--250.
- (317) Norderhaug, A.; Bakkevik, B. & Skogen, A. (1997), '*Dactylorhiza sambucina*, a threatened species in Norway?', *Blyttia* **55**(2), 73--86.
- (318) Norderhaug, A.; Bakkevik, B. & Skogen, A. (1997), '*Dactylorhiza sambucina*, a threatened species in Norway?' [norwegisch], *Blyttia* **55**(2), 73-86.
- (319) Oester, M.; Ask, K.; Roemermann, C.; Tackenberg, O. & Eriksson, O. (2009), 'Plant colonization of ex-arable fields from adjacent species-rich grasslands: The importance of dispersal vs. recruitment ability', *Agriculture Ecosystems & Environment* **130**(3-4), 93--99.
- (320) von Oheimb, G. & Brunet, J. (2007), 'Dalby Söderskog revisited: long-term vegetation changes in a south Swedish deciduous forest', *Acta Oecologica* **31**(2), 229 - 242.
- (321) Oklejewicz, K. (1997), '*Veronica gentianoides* (Scrophulariaceae): An interesting synanthropic species in the Polish flora', *Fragmenta Floristica et Geobotanica Series Polonica* **4**(0), 377--378.
- (322) Oldhammer, B. (1994), 'History of fires in central and northwestern Dalarna', *Svensk Botanisk Tidskrift* **88**(5), 259--266.
- (323) Olsson, K.-A. (2006), 'New endangered vascular plants in Skane', *Botaniska Notiser* **139**(3), 22-

33.

- (324) Onipchenko, V. G.; Blinnikov, M. S. & Semenova, G. V. (1994), 'Experimental investigation of alpine plant communities in the Northwestern Caucasus: Shading experiments in alpine grasslands', *Veroeffentlichungen des Geobotanischen Institutes der Eidgenoessische Technische Hochschule Stiftung Ruebel in Zuerich* **0**(115), 89--99.
- (325) Onipchenko, V. G.; Makarov, M. I. & van der Maarel, E. (2001), 'Influence of alpine plants on soil nutrient concentrations in a monoculture experiment', *Folia Geobotanica* **36**(3), 225--241.
- (326) Oostermeijer, J. G. B.; Luijten, S. H. & den Nijs, J. C. M. (2003), 'Integrating demographic and genetic approaches in plant conservation', *Biological Conservation* **113**(3), 389 - 398.
- (327) Oster, M.; Ask, K.; Cousins, S. A. O. & Eriksson, O. (2009), 'Dispersal and establishment limitation reduces the potential for successful restoration of semi-natural grassland communities on former arable fields', *Journal of Applied Ecology* **46**(6), 1266-1274.
- (328) PARKER, J. S. (1976), 'B-chromosome System of *Hypochoeris-maculata* .1. B-distribution, Meiotic Behavior and Inheritance', *Chromosoma* **59**(2), 167--177.
- (329) PARKER, J. S.; AINSWORTH, C. C. & TAYLOR, S. (1978), 'B-chromosome System of *Hypochoeris-maculata* .2. B-effects On Meiotic A-chromosome Behavior', *Chromosoma* **67**(2), 123--143.
- (330) PARKER, J. S.; TAYLOR, S. & AINSWORTH, C. C. (1982), 'The B-chromosome System of *Hypochoeris-maculata* .3. Variation In B-chromosome Transmission Rates', *Chromosoma* **85**(2), 299--310.
- (331) Pavlov, V. N.; Onipchenko, V. G.; Aksenova, A. A.; Volkova, E. V.; Zueva, O. I. & Makarov, M. I. (1998), 'The role of competition in Alpine plant communities (the Northwestern Causacus): An experimental approach', *Zhurnal Obshchei Biologii* **59**(5), 453--476.
- (332) Pedersen, H. A. (1997), 'Two unnoticed occurrences of *Selaginella selaginoides*', *Flora og Fauna* **103**(1), 17--18.
- (333) Pedersen, H. A. E. (2006), 'Systematics and evolution of the *Dactylorhiza romana/sambucina* polyploid complex (Orchidaceae)', *Botanical Journal of the Linnean Society* **152**(4), 405-434.
- (334) Pegg, G. & Vessey, J. (1973), 'Chitinase activity in *Lycopersicon esculentum* and its relationship to the in vivo lysis of *Verticillium albo-atrum* mycelium', *Physiological Plant Pathology* **3**(2), 207 - 222.
- (335) PEGTEL, D. M. (1988), 'Germination In Declining and Common Herbaceous Plant-populations Co-occurring In An Acid Peaty Heathland', *Acta Botanica Neerlandica* **37**(2), 215--223.
- (336) PEGTEL, D. M.; BLOM, C. & KROEZE, C. (1987), 'Aluminum and Manganese Affected Growth of Threatened Plant Species Occurring On Poorly Buffered Sandy Soils of Moderate Acidity', *International Botanical Congress Abstracts* **17**, 392.

- (337) Peintinger, M. & Bergamini, A. (2006), 'Community structure and diversity of bryophytes and vascular plants in abandoned fen meadows', *Plant Ecology* **185**(1), 1--17.
- (338) Pellegrino, G. & Bellusci, F. (2009), 'Molecular identification of mycorrhizal fungi in *Dactylorhiza sambucina* (Orchidaceae)', *Biologia* **64**, 893-897.
- (339) Pellegrino, G.; Bellusci, F. & Musacchio, A. (2008), 'Double floral mimicry and the magnet species effect in dimorphic co-flowering species, the deceptive orchid *Dactylorhiza sambucina* and rewarding *Viola aethnensis*', *Preslia* **80**(4), 411-422.
- (340) Pellegrino, G.; Bellusci, F. & Musacchio, A. (2005), 'Evidence of post-pollination barriers among three colour morphs of the deceptive orchid *Dactylorhiza sambucina* (L.) Soo', *Sexual Plant Reproduction* **18**(4), 179-185.
- (341) Pellegrino, G.; Caimi, D.; Noce, M. E. & Musacchio, A. (2005), 'Effects of local density and flower colour polymorphism on pollination and reproduction in the rewardless orchid *Dactylorhiza sambucina* (L.) Soo', *Plant Systematics and Evolution* **251**(2-4), 119-129.
- (342) Pellegrino, G.; Caputo, P.; Cozzolino, S.; Menale, B. & Musacchio, A. (2000), 'Molecular characterization of a hybrid zone between *Orchis mascula* and *O. pauciflora* in Southern Italy', *BIOLOGIA PLANTARUM* **43**(1), 13-18.
- (343) Pellegrino, G.; Cozzolino, S.; D'Emerico, S. & Grunanger, P. (2000), 'The taxonomic position of the controversial taxon *Orchis clandestina* (Orchidaceae): Karyomorphological and molecular analyses', *BOTANICA HELVETICA* **110**(2), 101-107.
- (344) Pellegrino, G.; D'Emerico, S.; Musacchio, A.; Scrugli, A. & Cozzolino, S. (2005), 'Confirmation of hybridization among sympatric insular populations of *Orchis mascula* and *O. provincialis*', *PLANT SYSTEMATICS AND EVOLUTION* **251**(2-4), 131-142.
- (345) Pellegrino, G. & Musacchio, A. (2006), 'Effects of defoliation on reproductive success in two orchids, *Serapias vomeracea* and *Dactylorhiza sambucina*', *Annales Botanici Fennici* **43**(2), 123-128.
- (346) Peterken, G. (1977), 'Habitat conservation priorities in British and European woodlands', *Biological Conservation* **11**(3), 223 - 236.
- (347) Peterken, G. (1974), 'A method for assessing woodland flora for conservation using indicator species', *Biological Conservation* **6**(4), 239 - 245.
- (348) Peterken, G. F. & Francis, J. L. (1999), 'Open spaces as habitats for vascular ground flora species in the woods of central Lincolnshire, UK', *Biological Conservation* **91**(1), 55 - 72.
- (349) Pettersson, G. & Nilsson, L. A. (1983), 'The Pollination Ecology of *Dactylorhiza-sambucina* On the Island of Stora Karlso Sweden In the Baltic', *Svensk Botanisk Tidskrift* **77**(2), 123-132.
- (350) Pfeifer, M.; Passalacqua, N. G.; Bartram, S.; Schatz, B.; Croce, A.; Carey, P. D.; Kraudelt, H. &

- Jeltsch, F. (2010), 'Conservation priorities differ at opposing species borders of a European orchid', *Biological Conservation* **143**(9), 2207 - 2220.
- (351) Philippi, G. (1970), 'Calamagrostis phragmitoides (Hartm.), das Purpureitgras im Schwarzwald und in den Vogesen', *Beitraege zur Naturkundlichen Forschung in Suedwestdeutschland* **29**(2), 107-110.
- (352) Phoenix, G. K.; Johnson, D.; Grime, J. P. & Booth, R. E. (2008), 'Sustaining ecosystem services in ancient limestone grassland: importance of major component plants and community composition', *Journal of Ecology* **96**(5), 894--902.
- (353) Piessens, K. & Hermy, M. (2006), 'Does the heathland flora in north-western Belgium show an extinction debt?', *Biological Conservation* **132**(3), 382 - 394.
- (354) Piqueray, J.; Bottin, G.; Delescaille, L.-M.; Bisteau, E.; Colinet, G. & Mahy, G. (2010), 'Rapid restoration of a species-rich ecosystem assessed from soil and vegetation indicators: The case of calcareous grasslands restored from forest stands', *Ecological Indicators In Press, Corrected Proof*, - .
- (355) Plackowski, R. (2009), 'PHYTOSOCIOLOGICAL COMPARATIVE ANALYSIS OF TWO LOCALITIES OF CAREX PULICARIS L. IN THE ?ÓD? PROVINCE (CENTRAL POLAND)', *TEKA Komisji Ochrony i Kszta?towania ?rodowiska Przyrodniczego* **6**, 223--227.
- (356) PLACKOWSKI, R. (1986), 'New Localities of Rare Species of Plants Near Radomsko In the Province of Piotrkow Trybunalski Central Poland', *Fragmenta Floristica et Geobotanica* **31-32**(1-2), 45--48.
- (357) Plowright, C. B. (1896), 'On the recent additions to our knowledge of the uredine[modifier letter high tone bar] and ustilagine[modifier letter high tone bar] with special reference to the british species', *Transactions of the British Mycological Society* **1**, 90 - 105.
- (358) Poschenrieder, C.; Bech, J.; Llugany, M.; Pace, A.; Fenes, E. & Barcelo, J. (2001), 'Copper in plant species in a copper gradient in Catalonia (North East Spain) and their potential for phytoremediation', *Plant and Soil* **230**(2), 247--256.
- (359) Poschlod, P. & WallisDeVries, M. F. (2002), 'The historical and socioeconomic perspective of calcareous grasslands--lessons from the distant and recent past', *Biological Conservation* **104**(3), 361 - 376.
- (360) Prekorsek, B. (1972), 'Contribution To the Problems of Pseudoseasonal Polymorphism of Gentianella-praecox', *Bioloski Vestnik*(20), 17-29.
- (361) Przywara, L.; Pirog, H.; Konieczny, R. & Jach, M. (1996), 'The use of tissue culture methods in interspecific hybridization within the genus Trifolium L.', *Acta Biologica Cracoviensia Series Botanica* **38**, 53--.
- (362) Punt, W. (1984), 'Index to taxa and terms', *Review of Palaeobotany and Palynology* **42**(1-4),

365 - 369.

- (363) Punt, W. (1984), 'Umbelliferae', *Review of Palaeobotany and Palynology* **42**(1-4), 155 - 363.
- (364) Punt, W. & Hoen, P. (2009), 'The Northwest European Pollen Flora, 70: Asteraceae – Asteroideae', *Review of Palaeobotany and Palynology* **157**(1-2), 22 - 183.
- (365) Punt, W. & Hoen, P. P. (1995), 'Caryophyllaceae', *Review of Palaeobotany and Palynology* **88**(1-4), 83 - 272.
- (366) Punt, W. & Nienhuis, W. (1976), 'Gentianaceae', *Review of Palaeobotany and Palynology* **21**(2), 89 - 123.
- (367) Punt, W.; Weenen, J. S. D. L. V. & Oostrum, W. A. P. V. (1974), 'Primulaceae', *Review of Palaeobotany and Palynology* **17**(3-4), 31 - 70.
- (368) Quinn, R.; Lawton, J.; Eversham, B. & Wood, S. (1994), 'The biogeography of scarce vascular plants in Britain with respect to habitat preference, dispersal ability and reproductive biology', *Biological Conservation* **70**(2), 149 - 157.
- (369) Rae, D.; Armbruster, W.; Edwards, M. & Svengård-Barre, M. (2006), 'Influence of microclimate and species interactions on the composition of plant and invertebrate communities in alpine northern Norway', *Acta Oecologica* **29**(3), 266 - 282.
- (370) Ramsbottom, J. (1922), 'Orchid Mycorrhiza: With plates II-VII', *Transactions of the British Mycological Society* **8**(1-2), 28 - 61, IN2-IN7.
- (371) Reinhammar, L. (1998), 'Systematics of *Pseudorchis albida* s.l. (Orchidaceae) in Europe and North America', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* **126**(4), 363-382.
- (372) Reinhammar, L. (1995), 'Evidence for two distinctive species of *Pseudorchis* (Orchidaceae) in Scandinavia.', *NORDIC JOURNAL OF BOTANY* **15**(5), 469-481.
- (373) Reinhammar, L. & Hedren, M. (1998), 'Allozyme differentiation between lowland and alpine populations of *Pseudorchis albida* s.lat. (Orchidaceae) in Sweden', *NORDIC JOURNAL OF BOTANY* **18**(1), 7-14.
- (374) Reinhammar, L.; Olsson, E. & Sormeland, E. (2002), 'Conservation biology of an endangered grassland plant species, *Pseudorchis albida*, with some references to the closely related alpine *P-straminea* (Orchidaceae)', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* **139**(1), 47-66.
- (375) Reinhammar, L.-G. & Hedren, M. (1998), 'Allozyme differentiation between lowland and alpine populations of *Pseudorchis albida* s.lat. (Orchidaceae) in Sweden', *Nordic Journal of Botany* **18**(1), 7--14.
- (376) Reinhammar, L.-G.; Olsson, E. G. A. & Sormeland, E. (2002), 'Conservation biology of an endangered grassland plant species, *Pseudorchis albida*, with some references to the closely

- related alpine *P. straminea* (Orchidaceae)', *Botanical Journal of the Linnean Society* **139**, 47-66.
- (377) Reintal, M.; Tali, K.; Haldna, M. & Kull, T. (2010-09-01), 'Habitat preferences as related to the prolonged dormancy of perennial herbs and ferns', *Plant Ecology* **210**(1), Springer Netherlands, 111--123.
- (378) Reiné, R.; Chocarro, C. & Fillat, F. (2004), 'Soil seed bank and management regimes of semi-natural mountain meadow communities', *Agriculture, Ecosystems & Environment* **104**(3), 567 - 575.
- (379) Rich, T.; Holyoak, D.; Margetts, L. & Murphy, R. (1997), 'Hybridisation between *Gentianella amarella* and *G. anglica*', *Watsonia* **21**, 313-325.
- (380) Rich, T. C. G. & Woodruff, E. R. (1996), 'Changes in the vascular plant floras of England and Scotland between 1930-1960 and 1987-1988: The BSBI Monitoring Scheme', *Biological Conservation* **75**(3), 217 - 229.
- (381) Richter, F. & Zoepfel, B. (2006), 'Symphenological species groups of mountain meadows in the Eastern Ore Mountains (Germany) based on cluster analysis', *Hercynia* **39**(1), 51--68.
- (382) Ricotta, C. (2005), 'Additive partitioning of Rao's quadratic diversity: a hierarchical approach', *Ecological Modelling* **183**(4), 365 - 371.
- (383) Ricotta, C.; Chiarucci, A. & Avena, G. (2004), 'Quantifying the effects of nutrient addition on community diversity of serpentine vegetation using parametric entropy of type [alpha]', *Acta Oecologica* **25**(1-2), 61 - 65.
- (384) Rigat, M.; Àngels Bonet, M.; Garcia, S.; Garnatje, T. & Vallès, J. (2007), 'Studies on pharmaceutical ethnobotany in the high river Ter valley (Pyrenees, Catalonia, Iberian Peninsula)', *Journal of Ethnopharmacology* **113**(2), 267 - 277.
- (385) Romero, M. T. & Rico, H. E. (1986), '*Swertia-perennis* New-record L. For the Central Mountain System Spain', *Anales del Jardín Botánico de Madrid* **43**(1), 177-178.
- (386) Roo-Zielinska, E. & Solon, J. (1997), 'Effect of Geographical Location on Species Composition, Vegetation Structure, Diversity and Phytoindicative Characteristics in Pine Forests', *Environmental Pollution* **98**(3), 347 - 360.
- (387) Roquet, C.; Saez, L.; Aldasoro, J. J.; Susanna, A.; Alarcon, M. L. & Garcia-Jacas, N. (2008), 'Natural delineation, molecular phylogeny and floral evolution in *campanula*', *Systematic Botany* **33**(1), 203--217.
- (388) Rosenthal, G. & Fink, S. (1996), '*Pedicularis palustris* L. in the Bremen area: Distribution, ecology, and causes of its decline', *Abhandlungen Naturwissenschaftlichen Verein zu Bremen* **43**(2), 429-447.
- (389) ROSSI, W.; CORRIAS, B.; ARDUINO, P.; CIANCHI, R. & BULLINI, L. (1992), 'GENE VARIATION AND

GENE FLOW IN ORCHIS-MORIO (ORCHIDACEAE) FROM ITALY', *PLANT SYSTEMATICS AND EVOLUTION* **179**(1-2), 43-58.

- (390) Rosén, E. & Bakker, J. P. (2005), 'Effects of agri-environment schemes on scrub clearance, livestock grazing and plant diversity in a low-intensity farming system on Öland, Sweden', *Basic and Applied Ecology* **6**(2), 195 - 204.
- (391) Ryan, M. J. & Smith, D. (2004), 'Fungal genetic resource centres and the genomic challenge', *Mycological Research* **108**(12), 1351 - 1362.
- (392) RYCHLEWSKA, U.; HODGSON, D.; HOLUB, M.; BUDESINSKY, M. & SMITALOVA, Z. (1985), 'ON TERPENES .288. THE STRUCTURE OF 2-OXO-8-ALPHA-ANGELOYLOXY-11-ALPHA-ACETOXY-5-BETA-H,6-ALPHA-H,7-ALPHA-H- GUAI-1(10),3-DIEN-6,12-OLIDE, A SESQUITERPENE LACTONE FROM LASERPITIUM-PRUTENICUM L REVISION OF THE STEREOSTRUCTURES OF NATIVE 2-OXOGUAI-1(10),3-DIEN-6,12-OLIDES FROM THE SPECIES OF THE UMBELLIFERAE FAMILYS', *COLLECTION OF CZECHOSLOVAK CHEMICAL COMMUNICATIONS* **50**(11), 2607-2624.
- (393) S. P. Carter, J. P. & Slingsby, D. R. (1987), 'Soil and Vegetation of the Keen of Hamar Serpentine, Shetland', *Journal of Ecology* **75**, No. 1, 21-42.
- (394) Salzmann, C. C.; Cozzolin, S. & Schiestl, F. P. (2007), 'Floral scent in food-deceptive orchids: Species specificity and sources of variability', *PLANT BIOLOGY* **9**(6), 720-729.
- (395) Santos-Gonzalez, J. C.; Finlay, R. D. & Tehler, A. (2007), 'Seasonal dynamics of arbuscular mycorrhizal fungal communities in roots in a seminatural grassland', *Applied and Environmental Microbiology* **73**(17), 5613--5623.
- (396) Santos-González, J. C. (2007), 'Diversity of Arbuscular Mycorrhizal Fungi in Grasslands and Arable Fields', PhD thesis, Swedish University of Agricultural Sciences Uppsala.
- (397) Schatz, B. (2006), 'Fine scale distribution of pollinator explains the occurrence of the natural orchid hybrid X *Orchis bergonii*', *ECOSCIENCE* **13**(1), 111-118.
- (398) Schiestl, F. (2005), 'On the success of a swindle: pollination by deception in orchids', *NATURWISSENSCHAFTEN* **92**(6), 255-264.
- (399) Schiestl, F. P.; Johnson, S. D. & Raguso, R. A. (2010), 'Floral evolution as a figment of the imagination of pollinators', *Trends in Ecology & Evolution* **25**(7), 382 - 383.
- (400) Schleuning, M. & Matthies, D. (2009), 'Habitat Change and Plant Demography: Assessing the Extinction Risk of a Formerly Common Grassland Perennial', *Conservation Biology* **23**(1), 174--183.
- (401) Schleuning, M.; Niggemann, M.; Becker, U. & Matthies, D. (2009), 'Negative effects of habitat degradation and fragmentation on the declining grassland plant *Trifolium montanum*', *Basic and Applied Ecology* **10**(1), 61 - 69.

- (402) Schlüter, P. M. & Schiestl, F. P. (2008), 'Molecular mechanisms of floral mimicry in orchids', *Trends in Plant Science* **13**(5), 228 - 235.
- (403) Schmidt, K. & Jensen, K. (2000), 'Genetic structure and AFLP variation of remnant populations in the rare plant *Pedicularis palustris* (Scrophulariaceae) and its relation to population size and reproductive components', *American Journal of Botany* **87**(5), 678-689.
- (404) Schmiede, R. (2004), 'Vegetationskundliche Analyse und naturschutzfachliche Bewertung ausgewählter Grünlandbiotope im Osterzgebirge', Master's thesis, Technische Universität Dresden Fakultät für Forst-, Geo- und Hydrowissenschaften Institut für Geographie Lehrstuhl für Allgemeine Physische Geographie.
- (405) Schneider, G. (2009), 'Busch-Nelke - *Dianthus seguieri* subsp. *glaber* Celak. Merkblatt Artenschutz 2', Technical report, Bayrisches Landesamt für Umwelt (LFU).
- (406) Schoenswetter, P. (2006), 'Molecular evidence for glacial refugia of mountain plants in the European Alps (vol 14, pg 3549, 2005)', *Molecular Ecology* **15**(2), 573--573.
- (407) Scholz, H. (1971), 'Gegenwärtige Kenntnisse über die Verbreitung der *Calamagrostis phragmitoides* Hartm. (Purpureitgras) in Mitteleuropa.', *Philippia* **1**, 85-90.
- (408) Schonswetter, P.; Tribsch, A.; Barfuss, M. & Niklfeld, H. (2002), 'Several Pleistocene refugia detected in the high alpine plant *Phyteuma globulariifolium* Sternb. & Hoppe (Campanulaceae) in the European Alps', *Molecular Ecology* **11**(12), 2637--2647.
- (409) Schonswetter, P.; Tribsch, A.; Stehlik, I. & Niklfeld, H. (2004), 'Glacial history of high alpine *Ranunculus glacialis* (Ranunculaceae) in the European Alps in a comparative phylogeographical context', *Biological Journal of the Linnean Society* **81**(2), 183--195.
- (410) Schulz, L. D. (1999), *Rote Liste Farn- und Samenpflanzen*, Landesamt für Umwelt und Geologie.
- (411) Schumacher, W. ; Helfrich, H.-P. K. H. K. C. L. C. M. A. S. K. K. S. B. J. (2007), 'Erfolgskontrolle des Vertragsnaturschutzes anhand der Populationsgrößen und -entwicklung seltener und gefährdeter Farn- und Blütenpflanzen.'(148), Technical report, Landwirtschaftliche Fakultät der Universität Bonn, 170 pp.
- (412) SCHUMACKER, R. (1978), 'Effect of Burning On Heaths and Semi Natural Meadows of the Elsenborn Plateau Liege Province Belgium Floristic Phytosociologic and Ecologic Aspects', *Gehu, Jean-marie (ed.). Documents Phytosociologiques, Vol. li. (phytosociological Documents, Vol. li.). (in Fr.). 467p. Illus. Maps. J. Cramer: Lehre, West Germany. Isbn 3-7682-1196-7, 409--420.*
- (413) SCHUMACKER, R.; DIRKSE, G. & DE ZUTTERE, P. (1974), 'Carex-dioica New-record and Carex-pulicaris In the Nature Reserve Lands of the Hohe Venn Mountains Province of Liege Belgium', *Lejeunia*(74), 1-21.
- (414) SCHWABE, A. (1990), 'Syndynamic Processes In Nardo-callunetea Communities Changes In

- Fallow Land After Renewed Cattle Grazing and Life History of *Arnica-montana* L', *Carolinea* **48**, 45--68.
- (415) Schwabe, A.; Kratochwil, A. & Pignatti, S. (2007), 'Plant indicator values of a high-phytodiversity country (Italy) and their evidence, exemplified for model areas with climatic gradients in the southern inner Alps', *Flora - Morphology, Distribution, Functional Ecology of Plants* **202**(5), 339 - 349.
- (416) Schwabe-Braun, A. (1990), 'Veränderungen in montanen Borstgrasrasen durch Düngung und Brachlegung: *Antennaria dioica* und *Vaccinium vitis-idaea* als Indikatoren', *Tuexenia* **10**, 295-310.
- (417) Schweingruber, F. H. (2007), 'Stem anatomy of Caryophyllaceae', *Flora - Morphology, Distribution, Functional Ecology of Plants* **202**(4), 281 - 292.
- (418) Sciama, D.; Augusto, L.; Dupouey, J. L.; Gonzalez, M. & Dominguez, C. M. (2009), 'Floristic and ecological differences between recent and ancient forests growing on non-acidic soils', *Forest Ecology and Management* **258**(5), 600--608.
- (419) Scopece, G.; Cozzolino, S.; Johnson, S. D. & Schiestl, F. P. (2010), 'Pollination Efficiency and the Evolution of Specialized Deceptive Pollination Systems', *AMERICAN NATURALIST* **175**(1), 98-105.
- (420) Scopece, G.; Juillet, N.; Mueller, A.; Schiestl, F. P. & Cozzolino, S. (2009), 'Pollinator attraction in *Anacamptis papilionacea* (Orchidaceae): a food or a sex promise?', *PLANT SPECIES BIOLOGY* **24**(2), 109-114.
- (421) Sharma, I. K.; Jones, D. L. & French, C. J. (2003), 'Unusually high genetic variability revealed through allozymic polymorphism of an endemic and endangered Australian orchid, *Pterostylis* aff. *picta* (Orchidaceae)', *Biochemical Systematics and Ecology* **31**(5), 513 - 526.
- (422) Shefferson, R. P. (2009), 'The evolutionary ecology of vegetative dormancy in mature herbaceous perennial plants', *Journal of Ecology* **97**(5), 1000--1009.
- (423) Shefferson, R. P. & Tali, K. (2007), 'Dormancy is associated with decreased adult survival in the burnt orchid, *Neotinea ustulata*', *JOURNAL OF ECOLOGY* **95**(1), 217-225.
- (424) Silvertown, J.; Wells, D. A.; Gillman, M.; Dodd, M. E.; Robertson, H. & Lakhani, K. H. (1994), 'Short-term effects and long-term after-effects of fertilizer application on the flowering population of green-winged orchid *Orchis morio*', *Biological Conservation* **69**(2), 191 - 197.
- (425) SIMACHEV, V. I. (1980), 'Biological Foundations of Protection of Rare Relict Species of the Higher Plants of Leningrad Oblast Russian-sfsr Ussr With Special Reference To *Pulsatilla-vernalis* *Viscaria-alpina* and *Oxytropis-sordida*', *Botanicheskii Zhurnal (St. Petersburg)* **65**(5), 725--737.
- (426) Singh, R.; Kachroo, M. & Agrawal, S. S. (2009), 'Standardization of *Centipeda minima*,

- Mimusops elengi, Orchis mascula and Paeonia emodii', *INDIAN JOURNAL OF TRADITIONAL KNOWLEDGE* **8**(3, Sp. Iss. SI), 338-345.
- (427) Skuhrava, M.; Skuhravy, V. & Hellrigl, K. (2002), 'Gall midge fauna (Cecidomyiidae, Diptera) of South Tyrol (2): Gall midges in the National Park Stilfser Joch and in Val Badia in the Dolomite Alps.', *Gredleriana* **2**, 103--136.
- (428) Smejkal, M. (1976), 'Calamagrostis-phragmitoides New-record For the Czechoslovakian Flora', *Preslia (Prague)* **48**(2), 124-130.
- (429) Smith, S. E. & Read, D. J. (1997), Orchid mycorrhizas'Mycorrhizal Symbiosis (Second Edition)', Academic Press, London, pp. 349 - 375.
- (430) Smithson, A. (2006), 'Pollinator limitation and inbreeding depression in orchid species with and without nectar rewards', *NEW PHYTOLOGIST* **169**(2), 419-430.
- (431) Smithson, A. (2002), 'The consequences of rewardlessness in orchids: Reward-supplementation experiments with Anacamptis morio (Orchidaceae)', *AMERICAN JOURNAL OF BOTANY* **89**(10), 1579-1587.
- (432) SMUL (2009), 'Postkarte Orchidee des Jahres Orchis mascula'.
- (433) Somerville, D. C. & Nicol, H. I. (2002), 'Mineral content of honeybee-collected pollen from southern New South Wales', *Australian Journal of Experimental Agriculture* **42**(8), 1131--1136.
- (434) Soudzilovskaia, N. A. & Onipchenko, V. G. (2005), 'Experimental investigation of fertilization and irrigation effects on an alpine heath, northwestern Caucasus, Russia', *Arctic Antarctic and Alpine Research* **37**(4), 602--610.
- (435) Soudzilovskaia, N. A.; Onipchenko, V. G.; Cornelissen, J. H. C. & Aerts, R. (2005), 'Biomass production, N : P ratio and nutrient limitation in a Caucasian alpine tundra plant community', *Journal of Vegetation Science* **16**(4), 399--406.
- (436) Spitzer, L.; Benes, J.; Dandova, J.; Jaskova, V. & Konvicka, M. (2009), 'The Large Blue butterfly, Phengaris [Maculinea] arion, as a conservation umbrella on a landscape scale: The case of the Czech Carpathians', *Ecological Indicators* **9**(6), 1056 - 1063.
- (437) Stadler, K.; Koch, M.; Bernhardt, K. G. & Greimler, J. (2010), 'Spatial arrangement and genetic structure in Gentianella aspera in a regional, local, and temporal context', *Plant Systematics and Evolution* **286**(1-2), 7-19.
- (438) Stambouli-Meziane, H.; Bouazza, M. & Thinon, M. (2009), 'La diversité floristique de la végétation psammophile de la région de Tlemcen (nord-ouest Algérie)', *Comptes Rendus Biologies* **332**(8), 711 - 719.
- (439) STANOVA, V. & GRULICH, V. (1993), 'Floristical and Phytogeographical Characteristic of Rudava River Alluvium', *Biologia* **48**(4), 407--410.

- (440) Steffen, K. (1956), 'Endomitosen Im Endosperm Von *Pedicularis Palustris* L', *Planta* **47**(6), 625-652.
- (441) Steffen, K. (1955), 'Kern und Nucleolenwachstum bei endomitotischer Polyploidisierung-Ein Beitrag zur karyologischen Anatomie von *Pedicularis palustris* L.', *Planta* **45**(4), 379-394.
- (442) STEINBRUCK, G.; SCHLEGEL, M.; DAHLSTROM, I. & ROTTGER, B. (1986), 'CHARACTERIZATION OF INTERSPECIFIC HYBRIDS BETWEEN ORCHIS-MASCULA AND ORCHIS-PALLENS (ORCHIDACEAE) BY ENZYME ELECTROPHORESIS', *PLANT SYSTEMATICS AND EVOLUTION* **153**(3-4), 229-241.
- (443) Stetsyuk, N. O. (1995), 'The floristic findings in the lower course of the Vorskla River', *Ukrayins'kyi Botanichnyi Zhurnal* **52**(5), 639--646.
- (444) Stoicovici, L. (1984), 'Interspecific Relationships of the Glacial Relics *Swertia-perennis* L and *Pedicularis-sceptrum-carolinum* L In A Romanian Fen', *Vegetatio* **56**(3), 139-145.
- (445) STRID, A. (1979), 'Contribution To the Flora of Mount Kajmakcalan Voras Oros Northern Greece', *Annales Musei Goulandris* **4**(RECD. 1979).
- (446) Sudova, R. (2009), 'Different growth response of five co-existing stoloniferous plant species to inoculation with native arbuscular mycorrhizal fungi', *Plant Ecology* **204**(1), 135--143.
- (447) Tackenberg, O. (2001), 'Methoden zur Bewertung gradueller Unterschiede des Ausbreitungspotentials von Pflanzenarten', PhD thesis, Philipps-Universität Marburg.
- (448) Tali, K. (2004), 'SPECIES STRUCTURE OF NEOTINEA USTULATA', PhD thesis, Chair of Applied Ecology, Institute of Botany and Ecology, University of Tartu, Estonia.
- (449) Tali, K.; Fay, M. & Bateman, R. (2006), 'Little genetic differentiation across Europe between early-flowering and late-flowering populations of the rapidly declining orchid *Neotinea ustulata*', *BIOLOGICAL JOURNAL OF THE LINNEAN SOCIETY* **87**(1), 13-25.
- (450) Tali, K.; Foley, M. J. Y. & Kull, T. (2004), '*Orchis ustulata* L.', *Journal of Ecology* **92**(1), 174--184.
- (451) Tali, K. & Kull, T. (2001), 'Highly variable flowering time in *Orchis ustulata* (Orchidaceae): consequences for population dynamics', *NORDIC JOURNAL OF BOTANY* **21**(5), 457-466.
- (452) Tauscher, L. (1978), 'Zum Vorkommen von *Carlina acaulis* L. subsp. *simplex* (W. et. K.) Arcang. auf dem Pohlberg bei Annaberg (Westerzgebirge).', *Gleditschia* **6**, 141-143.
- (453) Tekinsen, K. K. & Güner, A. (2010), 'Chemical composition and physicochemical properties of tubera salep produced from some Orchidaceae species', *Food Chemistry* **121**(2), 468 - 471.
- (454) Ter Borg, S. J.; Janse, A. & Kwak, M. M. (1980), 'Life Cycle Variation In *Pedicularis palustris* (Scrophulariaceae)', *Acta Botanica Neerlandica* **29**(5-6), 397-406.

- (455) Terrab, A.; Ángeles Ortiz, M.; Talavera, M.; Ariza, M. J.; del Carmen Moriana, M.; García-Castaño, J. L.; Tremetsberger, K.; Stuessy, T. F.; Baeza, C. M.; Urtubey, E.; de Fátima Ruas, C.; Casimiro-Soriguer, R.; Balao, F.; Gibbs, P. E. & Talavera, S. (2009), 'AFLP and breeding system studies indicate vicariance origin for scattered populations and enigmatic low fecundity in the Moroccan endemic *Hypochaeris angustifolia* (Asteraceae), sister taxon to all of the South American *Hypochaeris* species', *Molecular Phylogenetics and Evolution* **53**(1), 13 - 22.
- (456) Thompson, K. & Hodgson, J. G. (1996), 'More on the biogeography of scarce vascular plants', *Biological Conservation* **75**(3), 299 - 302.
- (457) TIRA, S.; GALEFFI, C. & DELLEMON.EM (1969), 'Flavonoids of *Antennaria Dioica* Gaertn - A Mnr Study', *Annali Di Chimica* **59**(3), 284--.
- (458) Topic, J. & Ilijanic, L. (2001), '*Carex pulcaris* L. (Cyperaceae), a new species in Croatian flora', *Natura Croatica* **10**(4), 423--426.
- (459) Trejgell, A.; Bednarek, M. & Tretyn, A. (2009), 'Micropropagation of *Carlina acaulis*', *Acta Biologica Cracoviensia* **51**, 97-103.
- (460) Trejgell, A.; Dabrowska, G. & Tretyn, A. (2009), 'In vitro regeneration of *Carlina acaulis* subsp simplex from seedling explants', *Acta Physiologiae Plantarum* **31**(3), 445-453.
- (461) Trejgell, A. & Tretyn, A. (2007), 'Analysis of flowering ability of regenerated *Carlina acaulis* subsp simplex plants', *ACTA AGROBOTANICA* **60**, 39-44.
- (462) Tribsch, A. & Schonswetter, P. (2003), 'Patterns of endemism and comparative phylogeography confirm palaeoenvironmental evidence for Pleistocene refugia in the Eastern Alps', *Taxon* **52**(3), 477--497.
- (463) TRINAJSTIC, I. (1972), 'A Contribution To the Phytosociological Typological Studies of the Field Vegetation of the Croatian Mountain Slopes', *Poljoprivredna Znanstvena Smotra*(28), 95--101.
- (464) Trist, P. (1981), '*Fritillaria meleagris* L.: Its survival and habitats in suffolk, England', *Biological Conservation* **20**(1), 5 - 14.
- (465) Trivedi, M. R.; Morecroft, M. D.; Berry, P. M. & Dawson, T. P. (2008), 'Potential effects of climate change on plant communities in three montane nature reserves in Scotland, UK', *Biological Conservation* **141**(6), 1665 - 1675.
- (466) Tsavkelova, E. A.; Bömke, C.; Netrusov, A. I.; Weiner, J. & Tudzynski, B. (2008), 'Production of gibberellic acids by an orchid-associated *Fusarium proliferatum* strain', *Fungal Genetics and Biology* **45**(10), 1393 - 1403.
- (467) Tsiftsis, S.; Tsiripidis, I.; Karagiannakidou, V. & Alifragis, D. (2008), 'Niche analysis and conservation of the orchids of east Macedonia (NE Greece)', *Acta Oecologica* **33**(1), 27 - 35.
- (468) Turland, N. J. (2006), 'Lectotypification of *Campanula saxatilis*, *Phyteuma pinnatum* and

- Verbascum arcturus, Linnaean names of three taxa endemic to Crete', *Willdenowia* **36**(1), 303--309.
- (469) von Ubisch, G. (1936), 'Genetic Studies on the Nature of Hermaphroditic Plants in *Antennaria Dioica* (L) Gaertn', *Genetics* **21**(3), 282--294.
- (470) Uotila, P. & Ahti, T. (2009), 'Additions to the vascular flora of the Berezovye Islands (Koiviston saaret), Karelian Isthmus, Russia', *Memoranda Societatis pro Fauna et Flora Fennica* **85**(2), 33--44.
- (471) Ur-Rahman, A. & Zaman, K. (1989), 'Medicinal plants with hypoglycemic activity', *Journal of Ethnopharmacology* **26**(1), 1 - 55.
- (472) URBANSKA, K. (1969), 'Embryological Investigations In *Antennaria Gaertn* .v. Micro- and Macrosporogenesis In Natural Hybrids Between A *Carpatica* (wahlenb) Bluff Etlfingerh and A *Dioica* (l) Gaertn', *Acta Biologica Cracoviensia Series Botanica* **12**(1-2), 21--.
- (473) URBANSKA, K. (1965), 'Badania Cytologiczne Nad Doswiadczalnymi Mieszancami Pomiedzy *Antennaria Carpatica* (wahlb) Bl Et Fing la *Dioica* (l) Gaertn - Cytological Studies In Experimental Hybrids Between *Antennaria Carpatica* (wahlb) Bl Et Fing and A *Dioica* (l) Gaertn', *Acta Biologica Cracoviensia Series Botanica* **8**(1), 51--.
- (474) Valletta, A.; Attorre, F.; Bruno, F. & Pasqua, G. (2008), 'In vitro asymbiotic germination of *Orchis mascula* L', *PLANT BIOSYSTEMS* **142**(3), 653-655.
- (475) Van Den Berg, L. J. L.; Dorland, E.; Vergeer, P.; Hart, M. A. C.; Bobbink, R. & Roelofs, J. G. M. (2005), 'Decline of acid-sensitive plant species in heathland can be attributed to ammonium toxicity in combination with low pH', *New Phytologist* **166**(2), 551--564.
- (476) Vanderpoorten, A.; Delescaille, L.-M. & Jacquemart, A.-L. (2004), 'The bryophyte layer in a calcareous grassland after a decade of contrasting mowing regimes', *Biological Conservation* **117**(1), 11 - 18.
- (477) Varga, S. & Kytöviita, M.-M. (2010), 'Interrelationships between mycorrhizal symbiosis, soil pH and plant sex modify the performance of *Antennaria dioica*', *Acta Oecologica* **36**(3), 291 - 298.
- (478) Varga, S. & Kytöviita, M.-M. (2008), 'Sex-specific responses to mycorrhiza in a dioecious species', *Am. J. Bot.* **95**(10), 1225-1232.
- (479) Vaz, A.; Kerbauy, G. & Figueiredo-Ribeiro, R. (1998), 'Changes in soluble carbohydrates and starch partitioning during vegetative bud formation from root tips of *Catasetum fimbriatum* (Orchidaceae)', *PLANT CELL TISSUE AND ORGAN CULTURE* **54**(2), 105-111.
- (480) VERA DE LA PUENTE, M. L. (1985), 'Chorology of Acidophilous Taxa In the Cantabrian Mountains Spain', *Revista de Biología de la Universidad de Oviedo* **3**, 71--84.
- (481) VETTER, J. (1995), 'Isoflavones In Different Parts of Common *Trifolium* Species', *Journal of*

Agricultural and Food Chemistry **43**(1), 106--108.

- (482) Viciani, D. & Moggi, G. (1996), 'Notes on the high altitude flora at Pratomagno (Eastern Tuscany) and considerations on the effects of the methane gas pipeline', *Webbia* **51**(1), 59--81.
- (483) Vilcinskas, E. & Dabkeviciene, G. (2009), 'Qualitative and quantitative characteristics of clover (*Trifolium* spp.) species in the first year of growing', *Zemdirbyste-agriculture* **96**(4), 170--180.
- (484) Vitko, K. R. (1981), 'Ecology of Cotton-grass *Eriophorum latifolium* (Cyperaceae) In the Moldavian-SSR', *Izvestiya Akademii Nauk Moldavskoi SSR Seriya Biologicheskikh i Khimicheskikh Nauk* **2**(2), 30-32.
- (485) VOLKOVICH, V. B. (1972), 'Sex Ratios and Growth Characteristics In Diclinous Individuals of *Antennaria dioica*', *Botanicheskii Zhurnal (St. Petersburg)* **57**(10), 1278--1286.
- (486) de Vos, P. (2010), 'European materia medica in historical texts: Longevity of a tradition and implications for future use', *Journal of Ethnopharmacology* **132**(1), 28 - 47.
- (487) Waesch, G. (2006), 'Seed bank studies from meadows in the Thuringian Forest (Germany) subjected to different cultivation regimes - can an entire grassland regenerate from a seed bank?', *Tuexenia*(26), 275--295.
- (488) Walentoswki, H. & Obermeier, E. (1992), 'Mountain meadows with *Dactylorhiza sambucina* (L.) Soo at the Brotjacklriegel in the Vorderer Bayerischer Wald (Bavaria)', *Tuexenia* **0**(12), 193-208.
- (489) WALKER, R. B.; CHOWDAPPA, P. & GESSEL, S. P. (1993), 'Major-element Deficiencies In *Casuarina equisetifolia*', *Fertilizer Research* **34**(2), 127--133.
- (490) Waller, D. M. (1979), 'Models of mast fruiting in trees', *Journal of Theoretical Biology* **80**(2), 223 - 232.
- (491) Wallin, L.; Svensson, B. M. & Lonn, M. (2009), 'Artificial Dispersal as a Restoration Tool in Meadows: Sowing or Planting?', *Restoration Ecology* **17**(2), 270--279.
- (492) Wallnofer, B. (2008), 'On the alleged presence of *Carex pulicaris* (Cyperaceae) in Croatia', *Acta Botanica Croatica* **67**(2), 245--246.
- (493) Warcup, J. & Talbot, P. (1966), 'Perfect states of some rhizoctonias', *Transactions of the British Mycological Society* **49**(3), 427 - 435.
- (494) Warth"/Gumpelstadt, N. „A. (2005), 'Das Brand-Knabenkraut (*Orchis ustulata* L.) – Orchidee des Jahres', *Mitteilungen aus dem Biosphärenreservat Rhön*. **10**, 20-21.
- (495) Wattez, J. R. (1974), 'Marsh Lousewort *Pedicularis palustris* ssp. *serotina* L In the North of France', *Naturalistes Belges* **55**(6), 241-257.
- (496) Weber, R. W. & Webster, J. (2001), 'Teaching techniques for mycology: 14. Mycorrhizal

- infection of orchid seedlings in the laboratory', *Mycologist* **15**(2), 55 - 59.
- (497) Weeda, E. J. & Keulen, S. M. A. (2007), 'Changes in the vegetation of the kathager beemden reserve, west of heerlen', *Natuurhistorisch Maandblad* **96**(2), 21--29.
- (498) Weiss-Schneeweiss, H.; Stuessy, T. F.; Siljak-Yakovlev, S.; Baeza, C. M. & Parker, J. (2003), 'Karyotype evolution in South American species of *Hypochaeris* (Asteraceae, Lactuceae)', *Plant Systematics and Evolution* **241**(3-4), 171--184.
- (499) WELK, E. (2001), 'Arealkundliche Analyse und Bewertung der Schutzrelevanz seltener und gefährdeter Gefäßpflanzen Deutschlands.', PhD thesis, Halle, Univ. Math.-Nat. Fak..
- (500) Wells, T.; Rothery, P.; Cox, R. & Bamford, S. (1998), 'Flowering dynamics of *Orchis morio* L. and *Herminium monorchis* (L.) R.Br. at two sites in eastern England', *BOTANICAL JOURNAL OF THE LINNEAN SOCIETY* **126**(1-2), 39-48.
- (501) WELLS, T. C. E. (1976), 'Biological Flora of British-isles: *Hypochaeris maculata* L. (*Achyrophorus maculatus* (L.) Scop.)', *Journal of Ecology* **64**(2), 757--774.
- (502) Wells, T. C. E.; Rothery, P.; Cox, R. & Bamford, S. (1998), 'Flowering dynamics of *Orchis morio* L. and *Herminium monorchis* (L.) R. Br. at two sites in eastern England', *Botanical Journal of the Linnean Society* **126**(1-2), 39 - 48.
- (503) Werner, K. (2002), 'Comments to the revised edition of "Exkursionsflora von Deutschland, Band 4 (Kritischer Band)". 3. To the nomenclature of some species and subspecies.', *Schlechtendalia*(8), 1--13.
- (504) Westergaard, M. (1958), 'The Mechanism of Sex Determination in Dioecious Flowering Plants', *Advances in Genetics* **9**, 217 - 281.
- (505) Wheeler, B. R. & Hutchings, M. J. (2002), '*Phyteuma spicatum* L.', *Journal of Ecology* **90**(3), 581-591.
- (506) Whipps, J.; Haselwandter, K.; McGee, E. & Lewis, D. (1982), 'Use of biochemical markers to determine growth, development and biomass of fungi in infected tissues, with particular reference to antagonistic and mutualistic biotrophs', *Transactions of the British Mycological Society* **79**(3), 385 - 400.
- (507) Wilhalm, T. (2007), 'Vascular plants', *Gredleriana* **7**, 428--431.
- (508) Wittig, B. (1996), 'Preserving and revitalization of vegetation from a headwater slope by grazing. First results', *Abhandlungen Naturwissenschaftlichen Verein zu Bremen* **43**(2), 309--316.
- (509) WITTIG, B. (1987), 'A Fountain Peat-plant Community With *Carex-pulicaris* Near Verden West Germany', *Abhandlungen Naturwissenschaftlichen Verein zu Bremen* **40**(4), 395--402.

- (510) Wittig, R.; Becker, U. & Nawrath, S. (2010), 'Grassland loss in the vicinity of a highly prospering metropolitan area from 1867/68 to 2000–The example of the Taunus (Hesse, Germany) and its Vorland', *Landscape and Urban Planning* **95**(4), 175 - 180.
- (511) Witting, B. (1998), 'Comparison of recent and historical records on Caricion fuscae (acid small-sedge fen) in the Elbe-Weser region', *Drosera* **98**(2), 85--102.
- (512) Wohlfahrt, G.; Bahn, M.; Horak, I.; Tappeiner, U. & Cernusca, A. (1998), 'A nitrogen sensitive model of leaf carbon dioxide and water vapour gas exchange: application to 13 key species from differently managed mountain grassland ecosystems', *Ecological Modelling* **113**(1-3), 179 - 199.
- (513) Wohlfahrt, G. M. & U. (2010), 'Lungen-Enzian – Gentiana pneumonanthe Sumpf-Läusekraut – Pedicularis palustris Berg-Wohlverleih – Arnica montana Beinbrech – Narthecium ossifragum Schutzwürdige Pflanzen des Moorrandes'(11), Technical report, Biologische Schutzgemeinschaft Hunte Weser-Ems (BSH) Naturschutzverband Niedersachsen (NVN), Ljkoportrjt.
- (514) Wood, C.; Pritchard, H. & Miller, A. (2000), 'Simultaneous preservation of orchid seed and its fungal symbiont using encapsulation-dehydration is dependent on moisture content and storage temperature', *CRYO-LETTERS* **21**(2), 125-136.
- (515) Wotavová, K.; Balounová, Z. & Kindlmann, P. (2004), 'Factors affecting persistence of terrestrial orchids in wet meadows and implications for their conservation in a changing agricultural landscape', *Biological Conservation* **118**(3), 271 - 279.
- (516) Yeo, M. J. M.; Blackstock, T. H. & Stevens, D. P. (1998), 'The use of phytosociological data in conservation assessment: a case study of lowland grasslands in mid Wales', *Biological Conservation* **86**(2), 125 - 138.
- (517) Yu, F.-H.; Krüsi, B.; Schütz, M.; Schneller, J. J. & Wildi, O. (2008), 'Plant communities affect the species-area relationship on Carex sempervirens tussocks', *Flora - Morphology, Distribution, Functional Ecology of Plants* **203**(3), 197 - 203.
- (518) Yuglichek, L. S. (2003), 'Rare central European species in the eastern part of male polissia', *Ukrayins'kyi Botanichnyi Zhurnal* **60**(1), 41--47.
- (519) ZALUSKI, T. (1978), 'Rare Plants of the Pilica Tributary Area Poland', *Acta Universitatis Nicolai Copernici Biologia* **22**, 133--136.
- (520) Zibareva, L.; Volodin, V.; Saatov, Z.; Savchenko, T.; Whiting, P.; Lafont, R. & Dinan, L. (2003), 'Distribution of phytoecdysteroids in the Caryophyllaceae', *Phytochemistry* **64**(2), 499 - 517.
- (521) Zobel, M.; Otsus, M.; Runk, K. & Liira, J. (2005), 'Can long-distance dispersal shape the local and regional species pool?', *Folia Geobotanica* **40**(1), 35--44.
- (522) Zolala, H. A.; Polevova, S. V. & Severova, E. E. (2009), 'The Palynomorphology of

Campanulaceae Juss.', *Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody Otdel Biologicheskii* **114**(Part 1), 39--54.

- (523) Öckinger, E.; Eriksson, A. K. & Smith, H. G. (2006), 'Effects of grassland abandonment, restoration and management on butterflies and vascular plants', *Biological Conservation* **133**(3), 291 - 300.
- (524) Öster, M. & Eriksson, O. (2007), 'Sex ratio mediated pollen limitation in the dioecious herb *Antennaria dioica*', *Ecoscience* **14**(3), 387-398.
- (525) Öster, M.; Persson, K. & Eriksson, O. (2008), 'Validation of plant diversity indicators in semi-natural grasslands', *Agriculture, Ecosystems & Environment* **125**(1-4), 65 - 72.
- (526) Öztürk, M.; Çelik, A.; Güvensen, A. & Hamzaoglu, E. (2008), 'Ecology of tertiary relict endemic *Liquidambar orientalis* Mill. forests', *Forest Ecology and Management* **256**(4), 510 - 518.
- (527) ŠPULEROVÁ, J. (2008), 'SUCCESSION CHANGES IN EXTENSIVELY USED AGRICULTURAL LAND', *Ekológia (Bratislava)* **27**(1), 54–64.
- (528) (2006), Arzneipflanzenporträts, in Siegfried Bäuml, ed., 'Heilpflanzenpraxis Heute (First Edition)', Urban & Fischer, Munich, pp. 49 - 460.
- (529) (2010), 'Sächsisches Naturschutzgesetz'.
- (530) (2009), 'Artenschutzsteckbrief Böhmischer Enzian (*Gentianella bohemica*)', BUND, Technical report, BUND.
- (531) (2008), 'SEB Main Meeting, Marseille 2008', *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology* **150**(3, Supplement 1), Abstracts of the Annual Main Meeting of the Society of Experimental Biology, 6th - 10th July 2008, Marseille, France, S1 - S40.
- (532) (1983), 'Abstracts of papers presented at the 20th Annual Meeting of the Society of Cryobiology August 30-September 2, 1983, University of Cambridge, Cambridge, United Kingdom Session 1—Symposium: Cryopreservation of organs', *Cryobiology* **20**(6), 698 - 748.