

## Exercises for “Quantum Phase Transitions”

Summer 26

DR. L. JANSSEN

Exercise 1 (24.04.26)

**1. Landau functional for a first-order phase transition**

(5 pts.)

Consider the free-energy density

$$f(\varphi) = \frac{a}{2}\varphi^2 + \frac{b}{4}\varphi^4 + \frac{c}{6}\varphi^6, \quad (1)$$

which depends on the real order parameter  $\varphi$ . The parameter  $a = a(T)$  depends on the temperature  $T$ , the coefficients  $b$  and  $c$  are temperature-independent constants, and  $b < 0$ ,  $c > 0$ .

- Determine the extrema of the functional  $f(\varphi)$  in Eq. (1). List all possibilities and sketch  $f(\varphi)$  in each case.
- Calculate the critical value  $a_c$  of the parameter  $a$  where the position  $\varphi_{\text{eq}}(a)$  of the global minimum of  $f(\varphi)$  changes discontinuously.
- Sketch the free energy  $f(\varphi_{\text{eq}})$  as a function of the parameter  $a$  in the vicinity of the phase transition. Why is it a first-order phase transition?

Hint: Expand  $f(\varphi_{\text{eq}})$  up to first order in  $\delta a = a - a_c$  around  $\delta a = 0$ .

**2. Phase diagram of a two-order-parameter system**

(5 pts.)

Consider a system with two real order parameters  $\varphi_1$  and  $\varphi_2$ , whose free-energy density is given by

$$f(\varphi_1, \varphi_2) = \frac{r}{2}(\varphi_1^2 + \varphi_2^2) - \frac{g}{2}(\varphi_1^2 - \varphi_2^2) + \frac{u}{4}(\varphi_1^4 + \varphi_2^4) + \frac{v}{2}\varphi_1^2\varphi_2^2, \quad (2)$$

where  $u, v > 0$ .

- Determine all extrema of the functional  $f(\varphi_1, \varphi_2)$  in Eq. (2). Which values are taken by  $\varphi_1^2$ ,  $\varphi_2^2$  at these extrema?
- Which conditions have to be posed on  $\varphi_1^2$  and  $\varphi_2^2$ ? Discuss which phases (i.e., configurations of  $\varphi_1$  and  $\varphi_2$ ) are physically reasonable in which areas of the  $(r, g)$  plane.
- In each case, determine the state with the lowest free energy as function of  $r$  and  $g$ . Distinguish between  $u^2 < v^2$  and  $u^2 > v^2$ .
- Sketch the phase diagram in the  $(r, g)$  plane for  $u^2 < v^2$  and  $u^2 > v^2$ , respectively. What are the orders of the different phase transitions?