

Exercises for “Quantum Phase Transitions” SS 18

DR. L. JANSSEN

Exercise 3 (for 18.05.18, 13:00)

1. Generating functional for noninteracting real bosons (4 points)

Consider the (discretized) field theory of a noninteracting real scalar boson field $\varphi \equiv (\varphi_k)_{k=1}^M$ with action

$$S[\varphi] = \sum_{k,l=1}^M \frac{1}{2} \varphi_k K_{kl} \varphi_l, \tag{1}$$

and positive definite symmetric and real matrix $K = K^T$ (“kernel”).

(a) Show that the connected n -point correlation functions

$$\langle \varphi_{l_1} \varphi_{l_2} \dots \varphi_{l_n} \rangle \equiv \frac{1}{Z[0]} \int \prod_{k=1}^M \frac{d\varphi_k}{\sqrt{2\pi}} \varphi_{l_1} \varphi_{l_2} \dots \varphi_{l_n} \exp(-S[\varphi]) \tag{2}$$

can be obtained from the *generating functional*

$$Z[h] = \int \prod_{k=1}^M \frac{d\varphi_k}{\sqrt{2\pi}} \exp\left(-S[\varphi] + \sum_{k=1}^M h_k \varphi_k\right) \tag{3}$$

via suitable derivatives with respect to the external source (“magnetic field”) h .

(b) Show that the generating functional for a noninteracting real scalar boson field theory can be computed in closed form as

$$Z[h] = (\det K)^{-1/2} \exp\left(\sum_{k,l=1}^M \frac{1}{2} h_k (K^{-1})_{kl} h_l\right). \tag{4}$$

(c) Use the above result to show that the propagator $G_{kl}^{(2)} \equiv \langle \varphi_k \varphi_l \rangle$ and the connected four-point function $G_{klmn}^{(4)} = \langle \varphi_k \varphi_l \varphi_m \varphi_n \rangle$ can be written as

$$G_{kl}^{(2)} = (K^{-1})_{kl} \quad \text{and} \quad G_{klmn}^{(4)} = G_{kl}^{(2)} G_{mn}^{(2)} + G_{km}^{(2)} G_{ln}^{(2)} + G_{kn}^{(2)} G_{lm}^{(2)}. \tag{5}$$

2. Partition function for complex bosons (1 point)

Use the result of problem 1 to show that the partition function $Z \equiv Z[0]$ for the theory of noninteracting complex boson fields Φ, Φ^* is

$$Z = \int \prod_{k=1}^M \frac{d\phi_k^* d\phi_k}{2\pi i} \exp\left(-\sum_{k,l=1}^M \phi_k^* K_{kl} \phi_l\right) = (\det K)^{-1}, \tag{6}$$

with positive definite Hermitian kernel $K = K^\dagger$.

please turn over!

3. Susceptibility exponent γ in the large- N limit (5+3* points)

Consider the partition function for the theory of N complex boson fields Φ_a and Φ_a^* , $a = 1, \dots, N$, interacting via an ultralocal two-body interaction,

$$Z = \int \prod_{a=1}^N \mathcal{D}\Phi_a^*(\vec{x}) \mathcal{D}\Phi_a(\vec{x}) e^{-S[\Phi^*, \Phi]} \quad (7)$$

with action

$$S[\Phi^*, \Phi] = \int d^d \vec{x} \left[\sum_{a=1}^N (|\nabla \Phi_a(\vec{x})|^2 + t |\Phi_a(\vec{x})|^2) + \frac{\lambda}{2N} \left(\sum_{a=1}^N |\Phi_a(\vec{x})|^2 \right)^2 \right]. \quad (8)$$

t is the tuning parameter for a classical phase transition distinguishing the disordered phase for $t > t_c$ from an ordered phase for $t < t_c$. λ denotes the quartic coupling.

(a) Show that the partition function can be written as

$$Z = \int \prod_{a=1}^N \mathcal{D}\Phi_a^*(\vec{x}) \mathcal{D}\Phi_a(\vec{x}) \mathcal{D}\sigma(\vec{x}) e^{-S_0[\Phi^*, \Phi] - \int d^d \vec{x} \left[\frac{N}{2\lambda} \sigma^2(\vec{x}) + i\sigma(\vec{x}) |\Phi(\vec{x})|^2 \right]}, \quad (9)$$

where we have introduced the order-parameter field $\sigma(\vec{x})$ which couples to $|\Phi(\vec{x})|^2 \equiv \sum_{a=1}^N |\Phi_a(\vec{x})|^2$ and S_0 denotes the Gaussian part of the action S . (This is the so-called *Hubbard-Stratonovich* transformation.)

(b) Integrate over all components Φ_a , $2 \leq a \leq N$, except the first one, to obtain an effective theory in Φ_1 and σ . Consider the limit $N \rightarrow \infty$, argue that the saddle-point approximation discussed in class becomes exact in this limit, and use it to compute the free energy density.

Hint:

$$\frac{f}{Nk_B T} = (t + \sigma) |\Phi_1|^2 - \frac{\sigma^2}{2\lambda} + \int \frac{d^d \vec{k}}{(2\pi)^d} \ln(k^2 + t + \sigma),$$

with the saddle-point conditions

$$(t + \sigma) \Phi_1 = 0 \quad \text{and} \quad \sigma = \lambda \int \frac{d^d \vec{k}}{(2\pi)^d} \frac{1}{k^2 + t + \sigma} + \lambda |\Phi_1|^2,$$

where we have assumed uniform fields at the saddle point, rotated $i\sigma \rightarrow \sigma$, and rescaled $\Phi_1/\sqrt{N} \rightarrow \Phi_1$.

(c) Show that the theory exhibits a phase transition for $d > 2$ at $t_c = -\lambda \int \frac{d^d \vec{k}}{(2\pi)^d} \frac{1}{k^2}$ and that the inverse susceptibility $\chi^{-1} \propto t + \sigma$ satisfies the implicit equation

$$(t + \sigma) \left(1 + \lambda \int \frac{d^d \vec{k}}{(2\pi)^d} \frac{1}{k^2(k^2 + t + \sigma)} \right) = t - t_c \quad (10)$$

for $t > t_c$. What happens for $d \leq 2$?

(d) Assume an ultraviolet cutoff Λ in the integral over wavevectors and compute the scaling form of the susceptibility in the critical region $t + \sigma \rightarrow 0$ for (i) $d > 4$, (ii) $d = 4$, and (iii) $2 < d < 4$. Compare with the predictions from Landau theory for the original model (7).

Hint: (i) $\chi \propto |t - t_c|^{-1}$, (ii) $\chi \propto \frac{\ln|t - t_c|}{|t - t_c|}$, (iii) $\chi \propto |t - t_c|^{-2/(d-2)}$.