

Exercises for “Quantum Phase Transitions” SS 18

DR. L. JANSSEN

Exercise 4 (for 08.06.18, 13:00)

1. Anisotropic perturbation to the $O(2)$ Wilson-Fisher fixed point (10 points)

Consider the $O(2)$ model for the two-component boson field $\phi = (\phi_1, \phi_2)$ (with ϕ_1, ϕ_2 being real scalars) in the presence of an anisotropic perturbation:

$$S = \int d^d x \left[\frac{1}{2}(\nabla\phi_1)^2 + \frac{1}{2}(\nabla\phi_2)^2 + \frac{r}{2}(\phi_1^2 + \phi_2^2) + \frac{u}{4!}(\phi_1^4 + \phi_2^4) + \frac{2v}{4!}\phi_1^2\phi_2^2 \right]. \quad (1)$$

For $v \neq u$ the continuous $O(2)$ rotational symmetry is explicitly broken, but a residual \mathbb{Z}_4 symmetry (fourfold rotations by integer multiples of $\pi/2$) remains intact.

- Classify all possible symmetry-allowed operators with respect to their scaling dimension. Are there any relevant or marginal operators near $d = 4$ dimensions that have been omitted in Eq. (1)?
- Show that the one-loop RG flow of the suitably rescaled couplings in $d = 4 - \epsilon$ can be written as:

$$\frac{dr}{d\ell} = 2r + \frac{1}{2}u + \frac{1}{6}v \quad (2)$$

$$\frac{du}{d\ell} = \epsilon u - \frac{3}{2}u^2 - \frac{1}{6}v^2 \quad (3)$$

$$\frac{dv}{d\ell} = \epsilon v - \frac{2}{3}v^2 - uv \quad (4)$$

- Show that these equations reduce to the expected flow equations of the $O(2)$ model in the limit $u = v$.
- Determine the linearized RG flow in the vicinity of the Wilson-Fisher fixed point at $r = r^*$ and $u = v = u^*$:

$$\frac{d\delta g_i}{d\ell} = \sum_{j=1}^3 B_{ij}\delta g_j + \mathcal{O}(\delta g^2), \quad \delta g_i \equiv g_i - g_i^*,$$

with the “stability matrix” $B_{ij} = \left. \frac{\partial(dg_i/d\ell)}{\partial g_j} \right|_{g=g^*}$ and $(g_i) \equiv (r, u, v)$. Is the \mathbb{Z}_4 anisotropy $\propto u - v$ relevant or irrelevant at the Wilson-Fisher fixed point? What is the corresponding eigenvalue of the stability matrix?