Supplementary Table 2. Regions throughout the rest of the brain that showed differential BOLD signal for R ("remember") and K ("know") responses regardless of type of stimuli. Talairach coordinates (x,y,z) for peak activation are given. 19 or more contiguous significant voxels at $P < 0.002$, uncorrected.

Region	Nnew > Enew	Enew > Nnew	ER > Enew	Enew > ER	EK > Enew	Enew > EK	NR > Nnew	New > NR	NK > Nnew	Nnew >NK
Cerebellum, R	(23–51,–20)		(11,–85,–26)							
Cingulate gyrus, L							(–7,–26,32)			(–22,–41,23)
Inferior frontal gyrus, L			(–46,15,12)				(–46,34,4)		(–47,41,12)	
Inferior parietal, L					(–52,–59,39)				(–37,–59,42)	
Inferior parietal, R					(48,–59,41)				(38,–48,44)	
Inferior temporal gyrus, R		(49,–62,–9)								
Middle frontal gyrus, L			(–49,15,35)		(–33,47,17)		(–40,–1,46) (–50,14,37)		(–49,16,37)	
Middle frontal gyrus, R					(40,36,24)		(48,47,–1)			
Middle occipital gyrus, R		(64,–44,5)	(32,–80,11)							
Middle temporal gyrus, L		(–60,–49,3)	(–57,–32,–6)				(–52,–59,24)			
Middle temporal gyrus, R							(64,–40,6)		(64,–40,4)	
Postcentral gyrus, L			(–47,–2,51)							
Precentral gyrus, L					(–44,–3,49)				(–53,16,8)	
Precentral gyrus, R					(41,–3,52)					
Precuneus, L							(–10,–68,29)		(–17,–70,30)	
Precuneus, R		(10,59,33)					(10,–68,36)		(5,–73,45)	
Putamen	(–19,18,–5)									
Superior frontal gyrus, L		(–6,46,44)			(–31,51,19)		(–10,14,57)		(–12,12,60)	
Superior frontal gyrus, R		(5,46,37)			(21,58,19)					
Supramarginal gyrus, R							(47,–48,30)			
Superior occipital gyrus,L									(–36,–80,33)	
Cuneus, L					(–6,–76,36)					